

Biocatalytic β -glucosylation/ β -galactosylation of Rebaudioside C by glycosynthases

Ye-Yang Yu¹, Si-Yu Zhang¹, Jia-Hui Sun¹, Yu-Yang Li¹, Yao-Yao Zhang¹, Ai-Min Lu², Li Liu¹ and Josef Voglmeir^{1*} 

¹ Glycomics and Glycan Bioengineering Research Center (GGBRC), College of Food Science and Technology, Nanjing Agricultural University, Nanjing 210095, PR China

² College of Sciences, Nanjing Agricultural University, Nanjing 210095, PR China

* Corresponding author, E-mail: josef.voglmeir@njau.edu.cn

Abstract

Extracts of *Stevia rebaudiana* Bertoni plant leaves contain a mixture of diterpene glycosides which possess high sweetness and are used as non-caloric sweeteners in foods and beverages. One of the most abundant of these steviol glycosides is Rebaudioside C (Reb C), which bears one esterified β -D-glucose and one glycosidically linked α -L-rhamnosyl-(1 $\rightarrow$ 2)-[β -D-glucosyl-(1 $\rightarrow$ 4)]-D-glucose trisaccharide. In this work, we isolated Reb C from commercial Stevia extracts using an orthogonal normal-phase $\times$ reversed-phase purification strategy. We then demonstrated that Reb C could be used as a donor substrate for enzymatic trans glycosylation reactions using chemically synthesized 1-deoxy-1-fluoro-D-glucose and 1-deoxy-1-fluoro-D-galactose donors and two engineered glycosynthases derived from *Agrobacterium* and *Streptomyces* β -glycosidases. This chemoenzymatic glucosylation/galactosylation strategy may be of use to extend the current repertoire of steviol glycosides analogues for analysis or the sensory evaluation of these novel steviol glycosides.

Citation: Yu YY, Zhang SY, Sun JH, Li YY, Zhang YY, et al. 2024. Biocatalytic β -glucosylation/ β -galactosylation of Rebaudioside C by glycosynthases. *Food Materials Research* 4: e008 <https://doi.org/10.48130/fmr-0023-0043>

Introduction

Steviol glycosides are a group of glycosylated diterpenes that can be isolated from the plant leaves of *Stevia rebaudiana* Bertoni ('sweet herb'), native to Paraguay and Brazil, where it is used to sweeten yerba mate tea^[1–3]. Besides South America, the *Stevia rebaudiana* plant is now also grown commercially in countries including China, Japan, the USA, Korea, India, and Vietnam^[4]. The mixture of extracted steviol glycosides possesses an overall 100–150-fold sweetness compared with sucrose^[5] and its current market value of US\$ ~700 M (2022) is expected to surpass a market volume of US\$ 1B in 2027 (www.imarcgroup.com/stevia-market). The main constituents of Stevia extracts are steviol glycosides, which are steviol diterpenes with a series of mono-, di-, and trisaccharide moieties (Fig. 1).

The carbohydrate heterogeneity of steviol glycosides isolated from Stevia leaves has also affected the overall sweetness of the extract, given that the sweetness of each type of glycoside varies significantly when compared to sucrose^[6]. For example, the most abundant steviol glycoside present in stevia leaves is stevioside, which possesses 210-fold the sweetness of sucrose, whereas other glycosides such as Rebaudioside A or Rebaudioside B have a slightly lower sweetness of 200 and 150, respectively. Another main steviol glycoside present in stevia leaves is Rebaudioside C (Reb C). Reb C possesses a relatively low sweetness of 30 and contains a bitter aftertaste, as few studies have been conducted on it, we believe that changing the glycosylation of Reb C is a strategy to increase its sweetness, enhance its application, and extend the range of steviol glycosides analogues.

Several genetic and chemoenzymatic approaches have been developed to improve the sweetness of steviol glycosides. The application of physical and chemical mutagenesis techniques as well as transgenic tools allowed the generation of Stevia plant variants with increased and/or altered levels of the individual types of steviol glycosides^[7–11]. Modifications of steviol glycosides were also described using metabolically engineered yeast and bacteria in whole-cell biotransformation^[12–16]. However, the majority of reported research on changing the glycosylation of steviol glycosides uses enzyme preparations was comprehensively reviewed^[17] and can be divided into the use of recombinant glucosyltransferases^[18–22], fructofuranosidases^[23–26], cyclodextrin glucanotransferases^[27–31], glucosidases^[32–36], and galactosidases^[37–39]. Interestingly, no attempts to perform biotransformation of steviol glycosides using glycosynthases have been reported yet.

Glycosynthases are mutant variants of glycosidases, in which a catalytically relevant nucleophile in the substrate binding site (usually aspartic acid or glutamic acid residues), is replaced with a small non-nucleophilic amino acid residue (commonly alanine or serine)^[40]. While these mutant variants are inactive in hydrolyzing the glycosidases' native carbohydrate substrates in the presence of water. However, in the presence of alcohols which possess good leaving groups (i.e., 1-fluoryl or 1-aryl sugar donors), glycosynthases can catalyze glycosidic linkages at high yields but are void of hydrolase activity. In this work, we chemically synthesized α -glucosyl and α -galactosyl fluorides as donor substrates and applied two previously described glycosynthases derived from *Streptomyces* sp. (StspBGlcE383A, UniProt ID A0A0N6ZTF1)^[41] from *Agrobacterium* sp. (AgtuBGlcE358S, UniProt ID P12614)^[42] to synthesize Reb C




Fig. 1 *Stevia rebaudiana* Bertonii plant and the structure of relevant steviol glycoside (Reb) variants.

bearing additional glucosylated and galactosylated moieties. Given that Reb C is one of the most abundant steviol glycosides present in *Stevia* extracts, but has a significantly lower sweetness compared to other Rebaudiosides, the addition of glucose or galactose moieties may be of use for increasing the overall sweetness of *Stevia* extracts.

Materials and methods

Materials

Pyridine, acetic anhydride ((CH₃CO)₂O), sodium sulfate (Na₂SO₄), magnesium sulfate (MgSO₄) and dichloromethane (CH₂Cl₂) were obtained from Sinopharm Chemical Reagent Co. (Shanghai, China). Ethyl acetate (EtOAc), petroleum ether and methanol (MeOH) were obtained from General-Reagent Co. (Shanghai, China). *Stevia* leaves extract, HF-pyridine and sodium methoxide were obtained from Macklin Co. (Shanghai, China). Thin-layer chromatography (TLC) was performed on Merck 60 F₂₅₄ HPTLC silica gel-coated aluminum sheets. Flash chromatography was performed with silica gel with a 300-400 mesh size. Carbohydrates were visualized using an orcinol staining solution (consisting of 40 mg orcinol monohydrate dissolved in 20 mL of aqueous H₂SO₄ (3.6 M)). The StspBGlCE383A and AgtuBGlCE358S glycosynthase genes were synthesized by Tsingke (Nanjing, China), and *E. coli* BL21 (Invitrogen, Shanghai, China) was used as an expression system for the production of the recombinant glycosynthases. Kanamycin and isopropyl-β-D-1-thiogalactopyranoside (IPTG) were purchased from Sigma-Aldrich (Shanghai, China). LCMS-grade acetonitrile and formic acid were purchased from Merck (Nanjing, China) and Aladdin Ltd. (Shanghai, China).

Expression and purification of StspBGlCE383A and AgtuBGlCE358S

StspBGlCE383A and AgtuBGlCE358S (Supplemental File S1) were subcloned into a pET-30a expression vector and transformed into *E. coli* BL21 (DE3) competent cells, then plated on LB agar supplemented with 50 mg/L kanamycin. A 5 mL LB medium containing 50 mg/L kanamycin was inoculated with a single colony and cultivated at 37 °C overnight. One mL of the cell suspension was then transferred into 400 mL LB medium (37 °C, 200 rpm), and grown in log phase until an OD₆₀₀ value of

0.6 was reached. The recombinant protein expression was then conducted for 16 h at 18 °C in the presence of 1 mM IPTG. The cells were then pelleted by centrifugation at 4,000 g for 15 min and resuspended in 10 mL lysis buffer (50 mM Tris, 100 mM NaCl, 1% Triton X-100 and 1 mM PMSF, pH = 8.0). The cell suspension was lysed by sonication for 20 min on ice and centrifuged at 12,000 g at 4 °C for 20 min. The cleared supernatant was purified by Ni²⁺-nitrilotriacetate (Ni²⁺NTA) agarose affinity chromatography using five column volumes of washing buffer (50 mM NaCl, 50 mM Tris, pH = 8.0) before the elution of the recombinant proteins with imidazole buffer (50 mM NaCl, 50 mM Tris, 500 mM imidazole, pH = 8.0). Fractions showing the high absorbance at 280 nm were collected and stored at -80 °C after the addition of 20% (v/v) glycerol. Samples of recombinant protein were analyzed by SDS-PAGE after Coomassie Brilliant G-250 staining.

Synthesis of glycosyl fluorides

1-F-glucose (**4a**) and 1-F-galactose (**4b**) were synthesized from glucose (**1a**) and galactose (**1b**) according to the acetylation/fluorination/deacetylation scheme shown in Fig. 2. In a first step, per-acetylated monosaccharides (**2a**, **2b**) were generated by using acetic anhydride/pyridine. The resulting O-acetylated sugars were converted into the corresponding glycosyl 1-fluorides using 70% hydrogen fluoride-pyridine (HF-Py). After purification, the acetylated glycosyl fluorides (**3a**, **3b**) were then deacetylated using sodium methoxide, yielding the desired products 1-F-glucose (**4a**) and 1-F-galactose (**4b**). Product formation was monitored by TLC and MALDI-ToF-MS analysis using a Bruker Autoflex Speed instrument (equipped with a 1000 Hz Smartbeam-II laser) using 6-aza-2-thiothymine as a crystallization matrix.

Glycosylation of Reb C and product analysis

Enzymatic glycosylation reactions were performed in volumes of 200 μL which consisted of 5 mM of Reb C, 50 mM of glycosyl fluoride **4a** or **4b**, and recombinant glycosynthases StspBGlCE383A or AgtuBGlCE358S (5 mg/mL) sodium phosphate buffer (100 mM, pH 7.0). After 24 h at 37 °C, the reactions were terminated by heating at 95 °C for 10 min.

Reactants were analyzed using a LC-ESI-MS method which was based on an RP-HPLC method for steviol glycosides recommended by the Joint FAO/WHO Expert Committee on Food




Fig. 2 Synthesis scheme for 1-F-glucose (4a) and 1-F-galactose (4b) from glucose (1a) and galactose (1b). Reagents and conditions: (i) pyridine, acetic anhydride; (ii) 70% hydrogen fluoride-pyridine, 0 °C; (iii) sodium methoxide, methanol.

Additives^[26]. The analytes were separated on a Shimadzu LCMS 8040 system (Shimadzu Corporation, Kyoto, Japan) using a Cosmosil reversed-phase C18 column (Nacalai Tesque Co., Kyoto, Japan, 250 mm × 4.6 mm; 5 μm pore size) with a flow rate at 0.5 mL/min. Water acidified with formic acid (0.1%v/v) and 100% acetonitrile were used as mobile phases for the gradual elution of the analytes (Supplemental Table S1). Analytes were detected by LC-ESI-MS analysis using scan mode detection in the m/z range between 500–2,000.

Results and discussion

Protein expression and purification

The expression and purification of StspBGlcE383A and AgtuBGlcE358S were evaluated by SDS-PAGE (Fig. 3). Protein bands corresponding to the expected molecular weight of the recombinant proteins were observed after induction by IPTG and could also be observed in the soluble elution fraction nickel-chelation affinity chromatography.

Synthesis of 1,2,3,4,6-penta-O-acetyl-glucose (2a) and 1,2,3,4,6-penta-O-acetyl-galactose (2b)

Two mmol glucose **1a** or galactose **1b** was added in 2 mL dry pyridine. After the addition of 1.5 mL acetic anhydride, the mixture was stirred in a round-bottom flask at room temperature for 4 h. TLC analysis (2:1 petroleum ether-ethyl acetate) and MALDI-ToF mass spectrometry were used to judge the completion of the per-acetylation reaction (Supplemental Fig. S1). After adding 80 mL of ice-cold water, compounds **2a** or **2b** were extracted in 60 mL dichloromethane. The organic layer was collected and washed with 80 mL water and 80 mL of saturated sodium hydrogen carbonate. After the addition of anhydrous sodium sulfate, the suspension was filtered and the resulting clear solution dried under reduced pressure (**2a**: 735 mg, 94%; **2b**: 711 mg, 91%).

Synthesis of 1-F-2,3,4,6-tetra-O-acetyl-glucose (3a) and 1-F-2,3,4,6-tetra-O-acetyl-galactose (3b)

The per acetylated monosaccharides **2a** and **2b** were dissolved in 6 mL 70% HF-Py in a 50 mL falcon tube at 0 °C, then gently stirred for 30 h. The completion of the reaction was judged by TLC analysis (2:1 petroleum ether-ethyl acetate) and MALDI-ToF mass spectrometry (Supplemental Fig. S2). The reaction was stopped by adding 20 mL of saturated aqueous sodium hydrogen carbonate. After adding 60 mL of dichloromethane, the organic phase was washed once with 80 mL saturated copper sulfate solution and washed twice with 80 mL saturated sodium chloride solution. After the addition of anhydrous magnesium sulfate, the suspension was filtered and the resulting clear solution dried under reduced pressure. The brown residue was purified by silica column chromatography (2:1 petroleum ether-ethyl acetate), and the acetylated glycosyl 1-fluorides **3a** and **3b** were obtained. Concentration under reduced pressure yielded a pale-yellow powder (**3a**: 154 mg, 21%; **3b**: 132 mg, 18%). These compounds were stored in methanol at a concentration of 200 mM.

Synthesis of 1-F-α-glucose (4a) and 1-F-α-galactose (4b)

Due to the instability of components **4a** and **4b**, the deprotection reactions were only performed in the aliquoted amounts required for the subsequent enzymatic reaction. Typically, 10 μL of acetylated glycosyl fluoride **3a** or **3b** (200 mM in methanol) were mixed with 980 μL of methanol and 10 μL of methanolic 0.3 M sodium methoxide. The progress of the deacetylation reaction was completed after 4 h at room temperature as judged by TLC (7:2:1 ethyl acetate-methanol-water) and MALDI-ToF mass spectrometry (Supplemental Fig. S3). Methanol was removed under reduced pressure to yield a pale-yellow powder, which was then dissolved in 10 μL of distilled water and immediately used as an acceptor for the glycosynthase reactions of Reb C.




Fig. 3 SDS-PAGE analysis of recombinant (a) StspBGlcE383A and (b) AgtuBGlcE358S protein expressed in *E. coli* BL21 (DE3) cells. M: Protein marker, Lane 1: cell suspension before induction, Lane 2: cell suspension after induction, Lane 3: supernatant of cell lysis, Lane 4: purified recombinant protein, Lane 5: precipitate of cell lysis.




Fig. 4 Extracted ion masses of Reb C and its enzymatically glycosylated reaction products. The analytes were separated using a reversed-phase C18 column and detected using a m/z range between 500–2,000.

Enzymatic Reb C glycosylation

Reb C is one of the main steviol glycoside present in Stevia leaves. Compared to other Rebaudiosides, it possesses a relatively low sweetness, therefore glycosidic decorations with glucose and galactose may be a way to modulate its sweetness. The glycosynthases StspBGlcE383A and AgtuBGlcE358S, which were previously used to glycosylate *para*-nitrophenyl glucoside in high yield in the presence of 1-F- α -galactose or 1-F- α -glucose^[41, 42], were also considered for this study. However, given that Reb C has three terminal glucose moieties, we anticipated that these biocatalysts could be also used for glycosylation reactions of this natural sweetener.

Initial enzymatic reaction trials using *para*-nitrophenyl- α - and β -glucosides as acceptor substrates and 1-F- α -glucose showed moderate glycosylation efficiency (between 8%–35% product formation, [Supplemental Fig. S4](#)) and encouraged us to apply Reb C as acceptor substrate. Using LC-ESI-MS in scan mode of Reb C ([Supplemental Fig. S5](#)) and the enzymatic reaction mixtures containing Reb C in the presence of 1-F- α -galactose or 1-F- α -glucose allowed the observation of additional mass signals corresponding to the addition of up to three extra galactose or glucose moieties on Reb C ([Fig. 4a–e](#)). Interestingly, StspBGlcE383A showed similar glycosylation abilities in reaction mixtures containing 1-F- α -galactose or 1-F- α -glucose, and allowed to add up to two hexose units to Reb C ([Fig. 4b, c](#)).

According to a previous study, StspBGlcE383A is able to form both β -1,3- and β -1,4-glycosylation, on the other hand, there are more than one hexose that can accept the glycosyl fluoride donor on Reb C, we believe this resulted in a diversity of glycosylated Reb C products. AgtuBGlcE358S was a more efficient biocatalyst and resulted in a quantitative glycosylation of the acceptor substrate (no observable Reb C mass signal present after the reaction) and allowed the addition of up to two glucose or up to three galactose units to Reb C ([Fig. 4d, e](#)). Although the feasibility of glycosylating Reb C with glycosynthases could be demonstrated, further optimization of this process may be required. By adding more enzyme to the reaction mixtures, higher glycosylation yields in shorter reaction times are expected as less glycoside fluoride donor is spontaneously degraded^[41, 43].

Conclusions

We demonstrated that chemically synthesized 1-deoxy-1-fluoro-D-glucose and 1-deoxy-1-fluoro-D-galactose could be used for the enzymatic transglycosylation of Reb C. The two applied glycosynthases were able to further glucosylate or galactosylate Reb C, and therefore allow the expansion of the current repertoire of steviol glycoside analogues. We envisage that both StspBGlcE383A and AgtuBGlcE358S may be also useful glycosynthases for glycosylating a variety of other natural products bearing terminal glucose moieties.

Author contributions

The authors confirm contribution to the paper as follows: study conception and design: Voglmeir J, Liu L; data collection: Yu YY, Zhang SY, Sun JH, Li YY, Zhang YY, Lu AM; analysis and interpretation of results: Voglmeir J, Liu L; draft manuscript preparation: Voglmeir J, Liu L. All authors reviewed the results and approved the final version of the manuscript.

Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgments

This work was supported in part by the National Natural Science Foundation of China (Grant Nos 31471703, 31671854, 31871793, and 31871754 to JV and LL), and the 100 Foreign Talents Plan (Grant No. JSB2014012 to JV).

Conflict of interest

The authors declare that they have no conflict of interest. Josef Voglmeir and Li Liu are the Editorial Board members of *Food Materials Research* who were blinded from reviewing or making decisions on the manuscript. The article was subject to the journal's standard procedures, with peer-review handled independently of these Editorial Board members and their research groups.

Supplementary information accompanies this paper at (<https://www.maxapress.com/article/doi/10.48130/fmr-0023-0043>)

Dates

Received 5 November 2023; Revised 17 December 2023; Accepted 27 December 2023; Published online 1 March 2024

References

1. Castro-Muñoz R, Correa-Delgado M, Córdova-Almeida R, Lara-Nava D, Chávez-Muñoz M, et al. 2022. Natural sweeteners: Sources, extraction and current uses in foods and food industries. *Food Chemistry* 370:130991
2. Shah Nawaz, Pandey DK, Konjengbam M, Dwivedi P, Kaur P, et al. 2021. Biotechnological interventions of in vitro propagation and production of valuable secondary metabolites in *Stevia rebaudiana*. *Applied Microbiology and Biotechnology* 105(23):8593–614
3. Srivastava V, Chaturvedi R. 2022. An interdisciplinary approach towards sustainable and higher steviol glycoside production from in vitro cultures of *Stevia rebaudiana*. *Journal of Biotechnology* 358:76–91
4. Kumari S, Gautam G, Sukany G, Meshram MR. 2019. Impact of spacing and levels of nitrogen on growth and yield of stevia (*Stevia rebaudiana* Bertoni). *Journal of Pharmacognosy and Phytotherapy* 8:1878–81
5. Cardello HMAB, Da Silva MAPA, Damasio MH. 1999. Measurement of the relative sweetness of stevia extract, aspartame and cyclamate/saccharin blend as compared to sucrose at different concentrations. *Plant Foods for Human Nutrition* 54(2):119–29
6. Prakash I, Markosyan A, Bunders C. 2014. Development of next generation Stevia sweetener: Rebaudioside M. *Foods* 3(1):162–75
7. Ahmad Khan S, Rahman LU, Verma R, Shanker K. 2015. Physical and chemical mutagenesis in *Stevia rebaudiana*: variant generation with higher UGT expression and glycosidic profile but with low photosynthetic capabilities. *Acta Physiologiae Plantarum* 38:4
8. Ahmad Khan S, Ur Rahman L, Shanker K, Singh M. 2014. Agrobacterium tumefaciens-mediated transgenic plant and somaclone production through direct and indirect regeneration from leaves in *Stevia rebaudiana* with their glycoside profile. *Protoplasma* 251(3):661–70

9. de Jesús Sánchez-Cordova Á, Capataz-Tafur J, Barrera-Figueroa BE, López-Torres A, Sanchez-Ocampo PM, et al. 2019. *Agrobacterium rhizogenes*-mediated transformation enhances steviol glycosides production and growth in *Stevia rebaudiana* plantlets. *Sugar Tech* 21(3):398–406
10. Wu Q, La Hovary C, Chen HY, Li X, Eng H, et al. 2020. An efficient *Stevia rebaudiana* transformation system and *in vitro* enzyme assays reveal novel insights into UGT76G1 function. *Scientific Reports* 10:3773
11. Zheng J, Zhuang Y, Mao HZ, Jang IC. 2019. Overexpression of *SrDXS1* and *SrKAH* enhances steviol glycosides content in transgenic *Stevia* plants. *BMC Plant Biology* 19:1
12. Li Y, Li Y, Wang Y, Chen L, Yan M, et al. 2016. Production of Rebaudioside A from Stevioside catalyzed by the engineered *Saccharomyces cerevisiae*. *Applied Biochemistry and Biotechnology* 178(8):1586–98
13. Wang Y, Chen L, Li Y, Li Y, Yan M, et al. 2016. Efficient enzymatic production of rebaudioside A from stevioside. *Bioscience, Biotechnology, and Biochemistry* 80:67–73
14. Shu W, Zheng H, Fu X, Zhen J, Tan M, et al. 2020. Enhanced heterologous production of glycosyltransferase UGT76G1 by co-expression of endogenous *prpD* and *malK* in *Escherichia coli* and its transglycosylation application in production of rebaudioside. *International Journal of Molecular Sciences* 21(16):5752
15. Gold ND, Fossati E, Hansen CC, DiFalco M, Douchin V, et al. 2018. A combinatorial approach to study cytochrome P450 enzymes for *de novo* production of steviol glucosides in Baker's yeast. *ACS Synthetic Biology* 7(12):2918–29
16. Moon JH, Lee K, Lee JH, Lee PC. 2020. Redesign and reconstruction of a steviol-biosynthetic pathway for enhanced production of steviol in *Escherichia coli*. *Microbial Cell Factories* 19:20
17. Gerwig GJ, Te Poele EM, Dijkhuizen L, Kamerling JP. 2016. *Stevia Glycosides: Chemical and enzymatic modifications of their carbohydrate moieties to improve the sweet-tasting quality*. In *Advances in Carbohydrate Chemistry and Biochemistry*, ed. Baker DC. Vol. 73. UK: Academic Press, Elsevier. pp. 1–72. <https://doi.org/10.1016/bs.accb.2016.05.001>
18. Chen L, Pan H, Cai R, Li Y, Jia H, et al. 2021. Bioconversion of Stevioside to Rebaudioside E Using Glycosyltransferase UGTSL2. *Applied Biochemistry and Biotechnology* 193(3):637–49
19. Dewitte G, Walmagh M, Diricks M, Lepak A, Gutmann A, et al. 2016. Screening of recombinant glycosyltransferases reveals the broad acceptor specificity of stevia UGT-76G1. *Journal of Biotechnology* 233:49–55
20. Ping Q, Yang L, Jiang J, Yuan J, Ai S, et al. 2022. Efficient synthesis of rebaudioside D2 through UGT94D1-catalyzed regio-selective glycosylation. *Carbohydrate Research* 522:108687
21. Yang L, Ping Q, Yuan Z, Jiang J, Guo B, et al. 2023. Highly efficient synthesis of mono- β -1,6-Glucosylated Rebaudioside A derivative catalyzed by glycosyltransferase YjIC. *Carbohydrate Research* 523:108737
22. Zhang J, Tang M, Chen Y, Ke D, Zhou J, et al. 2021. Catalytic flexibility of rice glycosyltransferase OsUGT91C1 for the production of palatable steviol glycosides. *Nature Communications* 12(1):7030
23. Yang Z, Uhler B, Zheng T, Adams KM. 1991. Transfructosylation of rebaudioside A (a sweet glycoside of *Stevia* leaves) with *Microbacterium* β -fructofuranosidase. *Chemical & Pharmaceutical Bulletin* 39(8):2043–45
24. Yang Z, B Uhler, T Zheng, KM Adams. 2019. Enzymatic synthesis and characterization of a novel α -1 $\rightarrow$ 6-glucosyl Rebaudioside C derivative sweetener. *Biomolecules* 9(1):27
25. Xu ZW, Li YQ, Wang YH, Bo Y, Ning ZX. 2009. Production of β -fructofuranosidase by *Arthrobacter* sp. and its application in the modification of Stevioside and Rebaudioside A. *Food Technology and Biotechnology* 47:137–43
26. Spohner SC, Czermak P. 2016. Enzymatic production of prebiotic fructo-oligosteviol glycosides. *Journal of Molecular Catalysis B: Enzymatic* 131:79–84
27. Prakash I, Chaturvedula VSP. 2014. Structures of some novel α -glucosyl diterpene glycosides from the glycosylation of Steviol glycosides. *Molecules* 19:20280–94
28. Wang W, Sun Y, Peng P, Gu G, Du G, et al. 2021. Two-step enzymatic conversion of Rebaudioside A into a Mono- α -1,4-Glucosylated Rebaudioside A Derivative. *Journal of Agricultural and Food Chemistry* 69:2522–30
29. Wellmann J, Wilms J, Hartmann B, Zirpel B, Brückner SI, et al. 2023. Novel α -glycosyl compounds from glycosylation of rubusoside. *Food Chemistry* 406:135033
30. Muñoz-Labrador A, Azcarate S, Lebrón-Aguilar R, Quintanilla-López JE, Galindo-Iranzo P, et al. 2020. Transglycosylation of steviol glycosides and Rebaudioside A: Synthesis Optimization structural analysis and sensory profiles. *Foods* 9:1753
31. Yu X, Yang J, Li B, Yuan H. 2015. High efficiency transformation of stevioside into a single mono-glycosylated product using a cyclodextrin glucanotransferase from *Paenibacillus* sp. CGMCC 5316. *World Journal of Microbiology and Biotechnology* 31:1983–991
32. Gerwig GJ, te Poele EM, Dijkhuizen L, Kamerling JP. 2017. Structural analysis of rebaudioside A derivatives obtained by *Lactobacillus reuteri* 180 glucanucrase-catalyzed trans- α -glucosylation. *Carbohydrate Research* 440-441:51–62
33. Yan Z, Cao X, Yang X, Yang S, Xu L, et al. 2021. A novel β -glucosidase from *Chryseobacterium scophthalmum* 1433 for efficient rubusoside production from stevioside. *Frontiers in Microbiology* 12:744914
34. Lan Q, Tang T, Yin Y, Qu X, Wang Z, et al. 2019. Highly specific sophorose β -glucosidase from *Sphingomonas elodea* ATCC 31461 for the efficient conversion of stevioside to rubusoside. *Food Chemistry* 295:563–68
35. Kang HJ, Lee HN, Hong SJ, Park BR, Ameer K, et al. 2022. Synthesis and characteristics of a rebaudioside-A like compound as a potential non-caloric natural sweetener by *Leuconostoc kimchii* dextranucrase. *Food Chemistry* 366:130623
36. Chaturvedula VSP, Upreti M, Prakash I. 2011. Structures of the novel α -glucosyl linked diterpene glycosides from *Stevia rebaudiana*. *Carbohydrate Research* 346(13):2034–38
37. Zerva A, Choroziou K, Kritikou AS, Thomaidis NS, Topakas E. 2021. β -glucosidase and β -galactosidase-Mediated transglycosylation of steviol glycosides utilizing industrial byproducts. *Frontiers in Bioengineering and Biotechnology* 9:685099
38. Wan HD, Xia YM. 2015. Enzymatic transformation of stevioside using a β -galactosidase from *Sulfolobus* sp. *Food & Function* 6(10):3291–95
39. Kitahata S, Ishikawa H, Miyata T, Tanaka O. 1989. Production of Rubusoside Derivatives by Transgalactosylation of Various α -Galactosidases. *Agricultural and Biological Chemistry* 53(11):2929–34
40. Williams SJ, Withers SG. 2002. Glycosynthases: Mutant glycosidases for glycoside synthesis. *Australian Journal of Chemistry* 55:3–12
41. aijes M, Saura-Valls M, Pérez X, Conti M, Planas A. 2006. Acceptor-dependent regioselectivity of glycosynthase reactions by *Streptomyces* E383A β -glucosidase. *Carbohydrate Research* 341(12):2055–65
42. Mayer C, Zechel DL, Reid SP, Warren RAJ, Withers SG. 2000. The E358S mutant of *Agrobacterium* sp. beta-glucosidase is a greatly improved glycosynthase. *FEBS Letters* 466(1):40–44
43. Li C, Zhu S, Ma C, Wang LX. 2017. Designer α 1,6-fucosidase mutants enable direct core fucosylation of intact N-glycopeptides and N-glycoproteins. *Journal of the American Chemical Society* 139(42):15074–87



Copyright: © 2024 by the author(s). Published by Maximum Academic Press on behalf of Nanjing Agricultural University. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.