

Genome-wide identification and expression profiling of the SWEET family in *Actinidia polygama* (Sieb. & Zucc.) Maxim.

Li Chen^{1#}, Hui-Fang Song^{2#}, Jia-Xin Liu^{1#}, Xu-Xin Jiang¹, Jun Ai¹, Zhen-Xing Wang^{1*} and Yun-Peng Wang^{3*}

¹ College of Horticulture, Jilin Agricultural University, Changchun, Jilin 130118, China

² Key Laboratory of Forest Silviculture and Conservation of the Ministry of Education, College of Forestry, Beijing Forestry University, Beijing 100083, China

³ Institute of Agricultural Biotechnology, Jilin Academy of Agricultural Sciences, Changchun 130033, China

Authors contributed equally: Li Chen, Hui-Fang Song, Jia-Xin Liu

* Corresponding authors, E-mail: zhenxinghd@aliyun.com; wangyunpengcas@163.com

Abstract

Sugar was transported from photosynthetic source cells to sink cells, sugar efflux transporter protein (sugars will eventually be exported to transporters, SWEETs) play an important role in the process. Although SWEET family members had been identified in many plants, transcriptome or genomics analysis of *Actinidia polygama* SWEET genes remains uncharacterized. In this study, 14 SWEET genes of *Actinidia polygama* were identified by protein Blast. The structural characteristics of SWEET genes showed that the number of amino acids encoded by the gene family was between 233 and 304, the relative molecular weight was between 25,918.83 and 33,192.12, the isoelectric point was within the range of 6.96 to 9.71, 14 ApSWEET from *Actinidia polygama* and the known grape and Arabidopsis SWEETs were divided into four clades (I, II, III, and IV) according to the phylogenetic relationships. The gene structure analysis showed that most of ApSWEET genes have six exons and five introns except ApSWEET5 and ApSWEET14. All ApSWEET proteins also contained P-loop, MtN3-slv, and transmembrane domain. Expression patterns of 14 ApSWEET in different organs and at different fruit developmental stages were analyzed. ApSWEET1 and ApSWEET5 exhibited tissue-specific expression, whereas other genes were more ubiquitously expressed. ApSWEET1, ApSWEET10, and ApSWEET11 exhibited higher expression in fruit. The results of this study provide insights into the characteristics of the SWEET genes in *Actinidia polygama* and may serve as a basis for further functional studies of such genes.

Citation: Chen L, Song HF, Liu JX, Jiang XX, Ai J, et al. 2024. Genome-wide identification and expression profiling of the SWEET family in *Actinidia polygama* (Sieb. & Zucc.) Maxim.. *Fruit Research* 4: e017 <https://doi.org/10.48130/frures-0024-0010>

Introduction

Sugars as the main soluble solid component are important nutrients and a key factor influencing the flavor quality of fruits^[1]. They also play a crucial role in regulating the expression of fruit-related genes, as well as plant growth and development^[2], stress responses^[3], and other developmental processes. In leaves, sugars are produced as important photo-assimilates and are loaded into the phloem system through the symplasm pathway. They are then unloaded in sink organs, such as fruits and flower, through the apoplast pathway^[4]. Sugar transporters that play an indispensable role in phloem loading, nectar secretion, and reproductive tissue development. Many studies have explored the mechanism of sucrose transport from source cells to sink cells, which involves the synergistic effects of multiple transporters^[4–8]. At present, sugar transporters in plants are classified into two types: Sugars Will Eventually be Exported Transporters (SWEETs) and major facilitator superfamily (MFS) transporters. Sugars Will Eventually be Exported Transporters (SWEETs) have been newly identified in plants in recent years^[9]. Sugar transporters of MFS are further classified into MSTs and SUTs^[10], which primarily facilitate sugar influx into the cytosol. However, some MSTs, namely the tonoplast sugar transporter (TST) and the vacuolar glucose transporter (VGT) involved in transporting sugars from the cytosol to vacuoles and act as H⁺/sugar antiporters^[11]. Both

MSTs and SUTs consist of 12 transmembrane α -helices and mediate membrane transport of different sugars^[12,13].

SWEET transporters differ from the classic 12 transmembrane structural domains of the major facilitator superfamily. The typical characteristics of SWEET genes include seven TM domains, including two MtN3_saliva domains, which are connected to a low conserved single TM, forming a 3-1-3 symmetric structure^[14]. Phylogenetic analysis shows that members of SWEET can be divided into four clades. Clades I, Clades II and Clades IV are mainly hexose transporters, while Clade III is mainly a sucrose transporter^[15]. There is substantial evidence suggesting that SWEET protein in Clade I and II may transport glucose, most of the Clade III SWEET proteins are sucrose transporters, and Clade IV evolved from other SWEET clades to primarily act as vacuolar transporters, regulating fructose transport^[14]. Members of the SWEET family are widely distributed, and SWEET genes are found not only in plants but also in prokaryotes and animals^[16]. With the development of plant genome research, the identification and functional study of SWEET family genes have been carried out in many plant species, and the number of SWEET family members varies significantly among different plants. For example, the SWEET family consists of 17 members in Arabidopsis, 22 in *Saccharum spontaneum*, 20 in strawberry, 15 in pomegranate, 19 in jujube, 29 in tomato, and 33 in apple^[17–23]. Previous studies have reported that members of the SWEET gene family are involved

in many important physiological processes of plant growth and development, including nectar production, seed, and pollen development, and the regulation of phloem loading, phloem transport, phloem unloading, abiotic stress, and pathogen interaction by regulating carbohydrate compounds^[18, 24–26]. SWEET proteins are also associated with flower, fruit, and seed development. For example, *AtSWEET8* mainly affected fertility during early inflorescence development, and *AtSWEET13* mainly affected fertility during late inflorescence development^[27]. The SWEET gene family was also involved in plant interactions with pathogens, such as *VvSWEET4*, which shows strong up-regulation of expression with infection of *Botrytis cinerea*^[28, 29].

Furthermore, SWEET proteins are key factor in regulating the distribution of soluble sugars, which is closely related to plant stress resistance^[26]. Overexpression of two homologues *AtSWEET16* and *AtSWEET17* in *Arabidopsis thaliana* can improve the cold resistance of transgenic plants^[26, 30]. Overexpression of *AtSWEET15* show more sensitivity to salt stress, and loss of function mutations in *AtSWEET15* show higher salt tolerance^[31]. Drought stress affects the redistribution of carbohydrates in plants^[32]. Under water deficiency conditions, the expression of *AtSWEET11* and *AtSWEET12* in leaves and roots increased, accompanying an increase in the transportation capacity of sucrose from leaves to roots, indicating that plants regulate the redistribution of carbohydrates by regulating the expression of SWEETs under water deficiency conditions^[33].

Actinidia polygama (Sieb. & Zucc.) Maxim. is a perennial vine plant, its fruit contains many unique and interesting flavors and many nutrients, including organic acids, amino acids, flavonoids, dietary fiber, and vitamins C and E. In addition, the leaves, fruits, stems, and roots also were exploited as nutraceuticals or medicine. Therefore, *A. polygama* as a valuable wild resource has very high nutritional and medicinal value. However, the current domestication of *A. polygama* is not complete, particularly in terms of improving fruit quality. During the ripening process of *A. polygama* fruit, sugar accumulation is closely related to fruit quality and edible ability. In this study, we identified the SWEET gene family members of *A. polygama*, and analyzed the physical and chemical properties, phylogenetic relationship, gene structure, and promoter elements of SWEET family proteins. We also studied the expression of SWEET gene family members in different organs of *A. polygama*, which laid a foundation for further study on the structure and function of the SWEET gene family.

Results

Identification and physical and chemical analysis of SWEET

By searching the SWEET gene on the website (ID: PRJDB13926), a total of 23 SWEET gene family members of *A. polygama* were screened out, and nine SWEET subfamily members were eliminated because of lack of the conserved structures. According to the physicochemical property table (Table 1), the amino acid quantity of the SWEET gene family ranged from 233 to 304. The molecular weight ranges from 25,895.58 to 33,192.12. The isoelectric point is between 6.96 and 9.71, and the stability coefficient of *ApSWEET4*, *ApSWEET5* and *ApSWEET14* proteins was greater than the remaining 11 proteins. The aliphatic index was 94.76~118.24, which was a stable protein. Furthermore, most of the SWEET genes were found to be present on the cell membrane, rarely on the chloroplast and Golgi apparatus.

Phylogenetic and structural analysis of SWEET gene family members in *A. polygama*

To study the phylogenetic relationships among SWEET genes in *A. polygama* and other plant species, a neighbor-joining phylogenetic tree was constructed by aligning 14 *ApSWEET* sequences, 17 *AtSWEET* sequences, and 14 *VvSWEET* sequences (Supplemental Tables S1–S3). Apparently, 45 proteins were clustered into four different groups (Fig. 1). In detail, six *ApSWEETs* (*ApSWEET9*, 10, 11, 12, 13, 14) showed high homology with three *AtSWEETs* (*AtSWEET1–3*) and three *VvSWEETs* (*VvSWEET2a*, 2b, 3) in group I. In group II, three *ApSWEETs* (*ApSWEET5*, 7, 8) were clustered with five *AtSWEETs* (*AtSWEET4–8*) and three *VvSWEETs* (*VvSWEET5a*, 5b, 7). *ApSWEET1/4* were homologous to seven *AtSWEETs* (9–15) and five *VvSWEETs* (*VvSWEET9*, 10, 11, 12, 15) in group III. Three *ApSWEET* (*ApSWEET2*, 3, 6), two *AtSWEETs* (*AtSWEET16*, 17) and three *VvSWEETs* (*VvSWEET17a*, 17b, 17d) were included in group IV. The exon-intron structural evolution showed that most of the *ApSWEET* contained six exons, except for *ApSWEET5* and *ApSWEET14* which contained five exons (Fig. 2). *ApSWEET2* and *ApSWEET4* had the shortest and longest sequence, respectively. *ApSWEET11*, *ApSWEET12*, and *ApSWEET13* had similar exon-intron structures. In addition, *ApSWEET1* and *ApSWEET4*, *ApSWEET3*, and *ApSWEET6* also showed similar structures, these genes belong to the same group. These results suggested that *ApSWEETs* in the same group shared similar exon-intron organizations.

Table 1. Physicochemical properties of SWEET gene family proteins.

Gene ID	Number of amino acids	Molecular weight	pI	Asp + Glu	Arg + Lys	Instability index	Predicted location
<i>ApSWEET1</i>	275	31,093	8.17	17	19	37.35 (stable)	Cell membrane
<i>ApSWEET2</i>	244	27,049.86	6.86	16	16	35.79 (stable)	Cell membrane
<i>ApSWEET3</i>	304	33,192.12	9.49	19	30	36.79 (stable)	Chloroplast
<i>ApSWEET4</i>	254	29,132.9	7.61	19	20	46.56 (unstable)	Cell membrane
<i>ApSWEET5</i>	259	28,602.2	9.71	13	24	45.53 (unstable)	Cell membrane
<i>ApSWEET6</i>	234	25,895.58	8.48	14	16	35.70 (stable)	Cell membrane
<i>ApSWEET7</i>	238	26,891.1	9.18	13	19	36.22 (stable)	Cell membrane
<i>ApSWEET8</i>	237	26,653.71	8.87	13	17	34.25 (stable)	Cell membrane
<i>ApSWEET9</i>	253	27,530.9	9.5	12	23	29.02 (stable)	Cell membrane Golgi apparatus
<i>ApSWEET10</i>	252	27,529.8	9.49	12	20	25.31 (stable)	Cell membrane
<i>ApSWEET11</i>	236	25,918.83	9.26	8	14	36.71 (stable)	Cell membrane
<i>ApSWEET12</i>	234	25,977.87	9.03	9	14	36.26 (stable)	Cell membrane
<i>ApSWEET13</i>	238	26,391.35	9.01	9	14	38.49 (stable)	Cell membrane
<i>ApSWEET14</i>	233	26,641.9	9.36	15	24	43.15 (unstable)	Cell membrane




Fig. 1 Phylogenetic analysis of the ApSWEETs from *A. polygama*, *Arabidopsis thaliana*, and *Vitis vinifera*. The Neighbor-joining tree was drawn using MEGA7.0 with 1,000 bootstraps. The roman numbers (I–IV) labeled with various colors indicate different clades: green – Clade I, orange – Clade II, red – Clade III, blue – Clade IV.

Analysis of conserved motif of ApSWEET gene family

The Motif1~Motif10 conserved motifs were found in the SWEET gene of *A. polygama* (Fig. 3), whereas motifs 1, 2, 4, and 5 were observed in all ApSWEET members. In addition, Motif3 was observed in 13 ApSWEET members except ApSWEET4. Six genes lacked Motif6, and five genes lacked Motif7. Only ApSWEET13, ApSWEET12 and ApSWEET2 genes contained Motif8, ApSWEET10, ApSWEET5 and ApSWEET4 contained Motif9, ApSWEET1 and ApSWEET14 contained Motif10. Most of the conserved motifs had a relatively consistent relationship with the evolutionary tree with the same order of number, suggesting that these genes had strong conserved structures and similar gene functions. We confirmed that the ApSWEET proteins also contained P-loop, MtN3-slv, and transmembrane domain (Fig. 4). The typical structure of plant SWEET proteins consists of seven predicted transmembrane (7-TM) helices forming two MtN3_slv domains (triple-helix bundles, THB) connected by a linker transmembrane helix (TM4). All ApSWEET genes comprise two sugar transporter domains for intercellular exchange.

Secondary and tertiary structure analysis of ApSWEET gene family members

SOPMA was used to analyze the secondary structure of the SWEET gene family in *A. polygama*, including Alpha helix, chain extension structure, Beta turn, and Random coil (Table 2). These results showed that Alpha helix and Random coil were significantly higher than the chain extension structure and Beta turn. In addition, the Alpha helix proportion of ApSWEET6 was the highest, and ApSWEET9 was the lowest. Meanwhile, the proportion of chain extension structure in ApSWEET9 was the highest, and ApSWEET3 was the lowest. The beta turn of ApSWEET8 was the highest, and ApSWEET4 was the lowest. A random coil of ApSWEET4 was the highest, but ApSWEET6 was the lowest.

SWISS-MODEL was used for homology modeling analysis (Fig. 5), and it was found that the three-dimensional structure of ApSWEET proteins could be roughly divided into two categories. ApSWEET1, ApSWEET2, ApSWEET4, ApSWEET5, ApSWEET6, ApSWEET7 and ApSWEET8 was clustered into one group, and ApSWEET3, ApSWEET9, ApSWEET10, ApSWEET11, ApSWEET12, ApSWEET13 and ApSWEET14 was clustered into one group.




Fig. 2 ApSWEET gene structure of *A. polygama*.




Fig. 3 The conserved motif analyses of ApSWEETs proteins.

Chromosomal localization and collinear analysis of the ApSWEET gene

Chromosome analysis of genes (Fig. 6) showed that except for chromosome ApChr19, other genes were evenly distributed on 11 chromosomes. Three ApSWEETs genes were distributed in clusters on chromosomes ApChr19, ApSWEET12, and ApSWEET13 may be due to gene replication. According to the collinearity analysis diagram (Fig. 7), ApSWEET2 and ApSWEET6, ApSWEET4 and ApSWEET7, ApSWEET11 and ApSWEET3/12/13 exist collinearity, which may be obtained by chromosome fragment replication.

Promoter analysis of the ApSWEET gene family

In order to study the potential regulatory factors of the ApSWEET gene, the 2,000 bp promoter region of this family was analyzed (Fig. 8), and 88 elements in promoter regions of all ApSWEETs genes were predicted. The results showed that

response elements such as low temperature, light, and hormone appeared in most gene promoter regions, indicating that genes may be affected by low temperature, light, and hormone levels. They were classified into three groups based on their functional associations: stresses (ARE, DRE, STRE, LTR, MBS, and MYC), hormones (ABRE, TATC-box, CGTCA motif/TGACG motif, HD-Zip1, P-Box, GARE-motif, GA-motif, ERE, and TCA-element) and light (GT1-motif, TCCC-motif, TCT-motif, G-Box, Gap-box, LAMP-element). Among these elements, six elements were responsive to stress, ten elements were responsive to hormones, and six elements were responsive to light. Four development-related elements are responsive to meristem expression (CAT-box), cis-regulatory element involved in endosperm expression (GCN4), involved in endosperm-specific negative expression (AACA), and seed-specific regulation (RY). These findings indicated that ApSWEETs may respond to hormones or be involved in plant growth and stress resistance.




Fig. 4 Conserved structural domains of ApSWEETs.

Table 2. Secondary structure analysis of ApSWEET family members.

	Alpha helix	Extended strand	Beta turn	Random coil
ApSWEET1	45.09%	16.73%	2.18%	36.00%
ApSWEET2	41.80%	22.54%	2.87%	32.79%
ApSWEET3	49.01%	15.46%	5.26%	30.26%
ApSWEET4	37.80%	18.11%	0.79%	43.31%
ApSWEET5	35.91%	21.24%	4.63%	38.22%
ApSWEET6	50.00%	21.79%	3.85%	24.36%
ApSWEET7	40.34%	21.85%	4.20%	33.61%
ApSWEET8	40.51%	21.94%	5.91%	31.65%
ApSWEET9	33.99%	25.30%	4.35%	36.36%
ApSWEET10	46.83%	17.06%	3.97%	32.14%
ApSWEET11	42.37%	19.49%	5.08%	33.05%
ApSWEET12	41.88%	20.94%	4.27%	32.91%
ApSWEET13	46.22%	19.33%	2.52%	31.93%
ApSWEET14	36.48%	21.89%	3.86%	37.77%

Expression analysis of ApSWEET gene

Real-time fluorescence PCR was used to detect the expression of ApSWEET members in leaves, stems, flowers, roots, mature fruits (Fig. 9) and fruits of different developmental stages (Fig. 10). These results showed that the expression level of ApSWEET2, ApSWEET3, ApSWEET10, ApSWEET11 and ApSWEET13 was higher in leaves, the expression level of ApSWEET2, ApSWEET3, ApSWEET10, ApSWEET11, ApSWEET9, and ApSWEET13 was higher in stem, the expression level of ApSWEET5 and ApSWEET11 was higher in flower, the expression level of ApSWEET3, ApSWEET10, and ApSWEET11 was higher in the root, the expression level of ApSWEET1, ApSWEET10 and ApSWEET11 was higher in fruit. Most of the SWEET genes were found to be ubiquitously expressed in all tissues except for ApSWEET1 and ApSWEET5, the two genes were specifically expressed in the fruit and flower.

During fruit development, only ApSWEET5 had higher expression at an early stage of fruit development, ApSWEET1, ApSWEET2, ApSWEET10, and ApSWEET11 had higher expression at the mid and late stages of fruit development (Fig. 10). In addition, the glucose content in *A. polygama* is higher than the fructose and sucrose content from the initial measurement on June 11th 2022 (Fig. 10). In the final measurement on September 27th 2022, the glucose content is 2.7 times and 5.4 times higher than the fructose content and sucrose content.

Therefore, glucose content was the highest, followed by fructose, and sucrose content was the lowest during *A. polygama* development.

After the mature fruits are harvested, only ApSWEET1 shows strong expression at different storage periods. ApSWEET10 showed strong expression only on the first day after harvest, but the expression levels of ApSWEET11 gradually decrease with prolonged fruit storage time (Fig. 11).

Discussion

Although SWEET genes have been extensively studied in various species, such as *Arabidopsis thaliana*^[17], *Hemerocallis fulva*^[34], *Chinese jujube*^[14], *Malus*^[35] and *Litchi chinensis*^[36], the presence of SWEET gene family in *A. polygama* has not been reported. In this study, we identified 14 members of the SWEET gene family in *A. polygama*. The number is less than three in *Arabidopsis* and seven in rice^[17, 18]. Gene duplication has been shown to contribute to the expansion of SWEET genes in soybean and potato, enabling them to adapt to environmental changes. This could explain why these species have more SWEET proteins than others. Moreover, different predictions of their physical and chemical properties suggest that SWEET genes may have diverse functions in plants^[17]. Eleven of the identified genes were located on different chromosomes, consistent with findings from *Arabidopsis*. It is speculated that different members of the gene family may perform specific biological functions in different plant tissues. However, further research is needed to elucidate their exact roles in plant growth and development.

The structure and number of conserved motifs among members of the SWEET gene family were similar, and there was also a relative relationship between the evolution tree and gene structure. For example, ApSWEET11, ApSWEET12 and ApSWEET13 exhibited similar gene structures and belonged to the same clade. However, some gene structures displayed noticeable differences, suggesting the occurrence of expansion, reduction, or mutation during the evolution of the SWEET gene family. These variations may be associated with the diverse functions of the genes. The SWEET family members generally consist of five to six exons in their structures. It is speculated that the diversity of gene function may arise from the loss or addition of exons during the evolutionary process of these family genes. Additionally, all the members of SWEET






Fig. 5 Single protein structure of ApSWEETs.

gene family in *A. polygama* contained four highly conserved motifs: Motif1, Motif2, Motif4 and Motif5. These conserved motifs may play a key role in the biological function of the SWEET protein of *A. polygama*.

The expression patterns in different organs are closely correlated with gene function and serve as a predictor of biological functions. Numerous studies have reported the involvement of SWEET genes in various physiological processes, which usually were associated with specific tissue expression patterns. Our results also revealed that some SWEET genes were ubiquitously expressed in the flower, fruit tissues, root, leaf and stem. For example, ApSWEET5 and ApSWEET11 were highly expressed in flower. Some studies have reported that SWEET genes may play an important role in reproductive development. For example, AtSWEET13 and AtSWEET14 were found to be expressed in the anther wall, responsible for facilitating sucrose efflux into locules to support pollen development and maturation^[37], and mutations in AtSWEET9 had been shown to impair nectar secretion^[38,39]. These genes were specifically expressed in

pollen. Similarly, OsSWEET11 and SbSWEET9-3 were highly expressed in the panicle^[40, 41], suggesting that these genes may be essential for reproduction. Additionally, ApSWEET1 and ApSWEET5 exhibited specific expression in fruit. Similarly, high expression of VvSWEET transporters in flowers and berries highlighted a putative important role in sugar partitioning during flower and fruit development^[4]. Furthermore, ApSWEET3 and ApSWEET10 displayed high expression in the roots (Fig. 5), while ApSWEET2, ApSWEET9, and ApSWEET10 were relatively highly expressed in the leaves and stem, similar to the function of AtSWEET17 as fructose transporter^[42], these genes were proposed to participate in the phloem loading of photoassimilate in leaves of *A. polygama*. Our results suggested these genes may be involved in flower development, as well as the short and long-distance transportation and distribution of sugars. However, the regulatory mechanisms underlying the expression of these genes require further clarification.

In fruit, sugar (such as glucose, sucrose and fructose) is an important index that determines the quality of fruit. Many






Fig. 8 The cis-elements in the promoter sequences of *ApSWEETs* gene in *A. polygama*.

studies have reported functions of *SWEET* genes in both sink and source organs, particularly in fruits like tomato, grape, apple and Chinese jujube. The different expression patterns of the *SWEET* gene during fruit development are closely related to their function and can be used to predict biological functions. For instance, in apples, the expression of *SWEET* genes at young (*MdSWEET1.1/2*, *MdSWEET2.4* and *MdSWEET3.5*) and ripe fruit development stages were different^[23]. In grape, the expression of *VvSWEET10*, *VvSWEET12*, and *VvSWEET15* was higher in young fruit, but *VvSWEET15* was more abundant in mature fruit^[28]. Furthermore, *SISWEET7a*, *SISWEET14* and *SISWEET15* in tomato fruit are responsible for the transportation of glucose, fructose and sucrose^[43], and *VvSWEET4* acted as a glucose transporter in grape^[28]. In our study, seven *ApSWEET* genes were expressed at five different fruit development stage. Among them, three *ApSWEET* genes exhibited high expression at the early stage of fruit development, and two *ApSWEET* genes showed high expression in ripe fruit. Moreover, there is a certain correlation between gene expression and homology, with some genes belonging to the same clade as *ApSWEET1*, *VvSWEET10*, *VvSWEET11* and *VvSWEET15*^[4, 23]. On the basis of sugar content during fruit development, it was also speculated that *ApSWEET1* and two *ApSWEET* (*ApSWEET2* and *ApSWEET3*) may be involved in the transportation and distribution of sucrose and fructose during fruit ripening, while *ApSWEET10* and *ApSWEET11* may play a role in transportation and distribution of glucose during fruit ripening.

Materials and methods

Plant materials

Sampled plants were grown outdoors on the campus of Jilin Agricultural University, Jilin, China (43°48'48" N, 125°24'15" E). The annual precipitation is 867 mm, and the annual highest and lowest temperatures are 35 and −40 °C, respectively. Each field plot was divided into three subplots, and the seedlings were planted in each plot (3 m × 4 m). The phosphorous (4.12 ± 0.31 g/Kg), nitrogen (32.17 ± 1.98 g/Kg) and potassium (3.51 ± 0.19 g/Kg) concentrations were sufficient. Seedlings in all experiments were of uniform size. The expression of *ApSWEET*

genes was detected in *A. polygama* has different tissues and fruits at different development stages. Leaf, flower, root, stem, and fruit were collected. Fruit collected at 10, 40, 60, 80, and 85 d after full bloom respectively. After the mature fruits are harvested, fruit were stored for 1, 2, 3, 4, and 5 d. All fresh plant samples were collected with three independent replicates and immediately frozen in liquid nitrogen, then stored at −80 °C.

Identification of gene family members

Actinidia polygama *SWEET* gene family was identified by protein Blast of the 17 Arabidopsis *SWEET* proteins against the *Actinidia polygama* genome database (<https://figshare.com/s/f46aea0009a54a6a0528>).

The NCBI CDD (www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi) and PFAM (<http://pfam.sanger.ac.uk/>) website were used to search for the conserved domains of the candidate members.

Physicochemical properties and structural analysis

For the protein sequences encoded by the gene family members obtained above, ExPASy (<http://web.expasy.org/>) was used to predict their molecular weight, isoelectric point, stability and other physicochemical properties, respectively. WoLFPSORT (www.genscript.com/wolf-psort.html) was used to predict protein subcellular localization, and TMHMM (<https://services.healthtech.dtu.dk/service.php?TMHMM-2.0>) used to predict transmembrane structure. The gene structure of *SWEET* gene family members of *Actinidia polygama* was analyzed according to the location information of introns and exons on chromosomes. MEME was used to predict the motif of the protein-conserved domain. The analysis results were visualized using TBtools and modified by AI. The online software SOPMA (http://npsa-pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_sopma.html) was used to predict the secondary structure of *SWEET* protein, including random curling, chain extension structure, alpha helix and beta-turn. Using the SWISS-MODEL to analyze homology modeling (www.swissmodel.expasy.org).

Construction of molecular evolutionary tree

The MUSCLE was used for sequence comparison of the gene family members and was beautified with GENEDOC. The




Fig. 9 ApSWEET expression of different tissues.

comparison results were clipped with trim AI and analyzed with IQ-TREE evolutionary tree. Finally, the exhibited (<http://tree.bio.ed.ac.uk/software/figtree/>) for the beautification of the illustration.

Chromosome localization and collinearity analysis of gene family

Based on the whole genome and location information downloaded from the *Actinidia polygama* genome database




Fig. 10 ApSWEET expression of different fruit development stage.

(<https://figshare.com/s/f46aea0009a54a6a0528>), TBtools was used to conduct chromosome localization, collinearity, and gene tandem repeat event analysis for all SWEET gene family members of *Actinidia polygama*, and the results were visualized. MCSan was used to compare the whole genome sequence of *Actinidia polygama*, and the collinearity relationship was

obtained. The homologous gene map was drawn with TB tools.

Gene family promoter analysis

The 2 kb nucleotide sequences upstream of the transcription starting points of 14 genes of the gene family were




Fig. 11 *ApSWEET* expression of different storage periods.

predicted using PlantCARE, and TBtools software was used for visualization.

RNA extraction and real-time PCR

To validate the reliability of RNA-Seq, qRT-PCR for transcripts was carried out as described by Yu et al.^[44]. Total RNA was extracted from leaves, flowers, root, stem, and fruit sampled simultaneously. For each experiment, 1 µg of clean RNA was

converted to cDNA using the PrimeScript™ RT reagent Kit (TaKaRa Bio., Dalian, China) according to the manufacturer's protocol. Gene-specific primers were designed using Primer5.0 software (Premier Biosoft). Gene expression was performed using the SYBR Green Real-time PCR kit (TaKaRa Bio). *ACTIN* was used as a housekeeping gene after examining its constitutive expression pattern from the RNA-seq results. Relative gene expression levels were calculated with the $2^{-\Delta\Delta C_T}$ method^[45].

The sequences of the primers used for qRT-PCR are listed in [Supplemental Table S4](#).

Author contributions

The authors confirm contribution to the paper as follows: study conception and design: Wang ZX, Wang YP; performing the research: Chen L, Song HF, Liu JX, Jiang XX, Ai J. All authors reviewed the results and approved the final version of the manuscript.

Data availability

All data generated or analyzed during this study are included in this published article and its supplementary information files.

Acknowledgments

This study was supported by the Natural Science Foundation of China (to ZXW, GLS and JA, 31870673) and Jilin Province Development and Reform Commission, Grant/Award (Number: 2022C037-1); Department of Science and Technology of Jilin Province, Grant/Award (Numbers: 20210204083YY, 202101013697JC). This work complies with Chinese law. We thank Professor Ya-dong Li from the Jilin Agricultural University, Changchun, for providing recommendations for data collection.

Conflict of interest

The authors declare that they have no conflict of interest.

Supplementary information accompanies this paper at (<https://www.maxapress.com/article/doi/10.48130/frures-0024-0010>)

Dates

Received 11 December 2023; Revised 10 February 2024; Accepted 17 February 2024; Published online 6 May 2024

References

- Patrick JW. 1997. Phloem unloading: sieve element unloading and post-sieve element transport. *Annual Review of Plant Biology* 48:191–222
- Hall AJ, Minchin PEH, Gould N, Clearwater MJ. 2017. A biophysical model of fruit development with distinct apoplasmic and symplasmic pathways. *Acta Horticulturae* 1160:367–374
- Xiao W, Sheen J, Jang JC. 2000. The role of hexokinase in plant sugar signal transduction and growth and development. *Plant Molecular Biology* 44:451–61
- Kühn C, Grof CPL. 2010. Sucrose transporters of higher plants. *Current Opinion in Plant Biology* 13:288–98
- White PJ, Ding G. 2023. Long-distance transport in the xylem and phloem. In *Marschner's Mineral Nutrition of Higher Plants*, 4th edition, eds. Rengel Z, Cakmak I, White PJ. Academic Press. pp. 73–104. <https://doi.org/10.1016/B978-0-12-819773-8.00002-2>
- Li Y, Liang G, Nai G, Lu S, Ma W, et al. 2023. *VaUS2* confers cold tolerance in transgenic tomato and *Arabidopsis* by regulation of sucrose metabolism and ROS homeostasis. *Plant Cell Reports* 42:505–20
- Balparda M, Bouzid M, Martinez MDP, Zheng K, Schwarzländer M, et al. 2023. Regulation of plant carbon assimilation metabolism by post-translational modifications. *The Plant Journal* 114:1059–79
- Garg V, Kühn C. 2022. Subcellular dynamics and protein-protein interactions of plant sucrose transporters. *Journal of Plant Physiology* 273:153696
- Chen LQ, Qu XQ, Hou BH, Sosso D, Osorio S, et al. 2011. Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. *Science* 335:207–11
- Bai Q, Chen X, Zheng Z, Feng J, Zhang Y, et al. 2023. Vacuolar Phosphate Transporter1 (VPT1) may transport sugar in response to soluble sugar status of grape fruits. *Horticulture Research* 10:uhac260
- Hedrich R, Sauer N, Neuhaus HE. 2015. Sugar transport across the plant vacuolar membrane: nature and regulation of carrier proteins. *Current Opinion in Plant Biology* 25:63–70
- Gautam T, Dutta M, Jaiswal V, Zinta G, Gahlaut V, et al. 2022. Emerging roles of SWEET sugar transporters in plant development and abiotic stress responses. *Cells* 11:1303
- Salvi P, Agarwal R, Kaja, Gandass N, Manna M, et al. 2022. Sugar transporters and their molecular tradeoffs during abiotic stress responses in plants. *Physiologia Plantarum* 174:e13652
- Yang C, Zhao X, Luo Z, Wang L, Liu M. 2023. Genome-wide identification and expression profile analysis of SWEET genes in Chinese jujube. *PeerJ* 11:e14704
- Gao Y, Wang ZY, Kumar V, Xu XF, Yuan DP, et al. 2018. Genome-wide identification of the SWEET gene family in wheat. *Gene* 642:284–92
- Kumawat S, Sharma Y, Vats S, Sudhakaran S, Sharma S, et al. 2022. Understanding the role of SWEET genes in fruit development and abiotic stress in pomegranate (*Punica granatum* L.). *Molecular Biology Reports* 49:1329–39
- Hir RL, Spinner L, Klemens PAW, Chakraborti D, de Marco F, et al. 2015. Disruption of the sugar transporters *AtSWEET11* and *AtSWEET12* affects vascular development and freezing tolerance in *Arabidopsis*. *Molecular Plant* 8:1687–90
- Hu W, Hua X, Zhang Q, Wang J, Shen Q, et al. 2018. New insights into the evolution and functional divergence of the SWEET family in *Saccharum* based on comparative genomics. *BMC Plant Biology* 18:270
- Liu HT, Lyu WY, Tian SH, Zou XH, Zhang LQ, et al. 2019. The SWEET family genes in strawberry: identification and expression profiling during fruit development. *South African Journal of Botany* 125:176–87
- Miao H, Sun P, Liu Q, Miao Y, Liu J, et al. 2017. Genome-wide analyses of SWEET family proteins reveal involvement in fruit development and abiotic/biotic stress responses in banana. *Scientific Reports* 7:3536
- Li Y, Feng S, Ma S, Sui X, Zhang Z. 2017. Spatiotemporal expression and substrate specificity analysis of the cucumber SWEET gene family. *Frontiers in Plant Science* 8:1855
- Feng CY, Han JX, Han XX, Jiang J. 2015. Genome-wide identification, phylogeny, and expression analysis of the SWEET gene family in tomato. *Gene* 573:261–72
- Wei X, Liu F, Chen C, Ma F, Li M. 2014. The *Malus domestica* sugar transporter gene family: identifications based on genome and expression profiling related to the accumulation of fruit sugars. *Frontiers in Plant Science* 5:569
- Eckardt NA. 2021. Sweeter than SWEET: a single-cell leaf vasculature transcriptome atlas. *The Plant Cell* 33:445–46
- Chen LQ. 2014. SWEET sugar transporters for phloem transport and pathogen nutrition. *New Phytologist* 201:1150–55
- Chandran D. 2015. Co-option of developmentally regulated plant SWEET transporters for pathogen nutrition and abiotic stress tolerance. *IUBMB Life* 67:461–71
- Guan Y, Huang X, Zhu J, Gao J, Zhang H, et al. 2008. RUPTURED POLLEN GRAIN1, a member of the MtN3/saliva gene family, is

- crucial for exine pattern formation and cell integrity of microspores in *Arabidopsis*. *Plant Physiology* 147:852–63
28. Chong J, Piron MC, Meyer S, et al. 2014. The SWEET family of sugar transporters in grapevine: VvSWEET4 is involved in the interaction with *Botrytis cinerea*. *Journal of Experimental Botany* 65:6589–601
 29. Breia R, Conde A, Pimentel D, Conde C, Fortes AM, et al. 2020. VvSWEET7 is a mono- and disaccharide transporter up-regulated in response to *botrytis cinerea* infection in grape berries. *Frontiers in Plant Science* 10:1753
 30. Klemens PAW, Patzke K, Deitmer J, Spinner L, Le Hir R, et al. 2013. Overexpression of the vacuolar sugar carrier AtSWEET16 modifies germination, growth, and stress tolerance in *Arabidopsis*. *Plant Physiology* 163:1338–52
 31. Seo PJ, Park JM, Kang SK, Kim SG, Park CM. 2011. An *Arabidopsis* senescence-associated protein SAG29 regulates cell viability under high salinity. *Planta* 233:189–200
 32. Julius BT, Leach KA, Tran TM, Mertz RA, Braun DM. 2017. Sugar transporters in plants: new insights and discoveries. *Plant and Cell Physiology* 58:1442–60
 33. Durand M, Porcheron B, Hennion N, Maurousset L, Lemoine R, et al. 2016. Water deficit enhances C export to the roots in *Arabidopsis thaliana* plants with contribution of sucrose transporters in both shoot and roots. *Plant Physiology* 170:1460–79
 34. Huang D, Chen Y, Liu X, Ni D, Bai L, et al. 2022. Genome-wide identification and expression analysis of the SWEET gene family in daylily (*Hemerocallis fulva*) and functional analysis of HfSWEET17 in response to cold stress. *BMC Plant Biology* 22:211
 35. Nie P, Xu G, Yu B, Lyu D, Xue X, et al. 2022. Genome-wide identification and expression profiling reveal the potential functions of the SWEET gene family during the sink organ development period in apple (*Malus × domestica* Borkh.). *Agronomy* 12:1747
 36. Xie H, Wang D, Qin Y, Ma A, Fu J, et al. 2019. Genome-wide identification and expression analysis of SWEET gene family in *Litchi chinensis* reveal the involvement of LcSWEET2a/3b in early seed development. *BMC Plant Biology* 19:499
 37. Wang J, Xue X, Zeng H, Li J, Chen L. 2022. Sucrose rather than GA transported by AtSWEET13 and AtSWEET14 supports pollen fitness at late anther development stages. *New Phytologist* 236:525–37
 38. Schmitt AJ, Roy R, Klinkenberg PM, Jia M, Carter CJ. 2018. The octadecanoid pathway, but not COI1, is required for nectar secretion in *Arabidopsis thaliana*. *Frontiers in Plant Science* 9:1060
 39. Lin IW, Sosso D, Chen LQ, Gase K, Kim SG, et al. 2014. Nectar secretion requires sucrose phosphate synthases and the sugar transporter SWEET9. *Nature* 508:546–49
 40. Fei H, Yang Z, Lu Q, Wen X, Zhang Y, et al. 2021. OsSWEET14 cooperates with OsSWEET11 to contribute to grain filling in rice. *Plant Science* 306:110851
 41. Xiao Q, Zhen L, Wang Y, Hou X, Wei X, et al. 2022. Genome-wide identification, expression and functional analysis of sugar transporters in sorghum (*Sorghum bicolor* L.). *Journal of Integrative Agriculture* 21:2848–64
 42. Eom JS, Chen LQ, Sosso D, Julius BT, Lin IW, et al. 2015. SWEETs, transporters for intracellular and intercellular sugar translocation. *Current Opinion in Plant Biology* 25:53–62
 43. Zhang X, Feng C, Wang M, Li T, Liu X, et al. 2021. Plasma membrane-localized SISWEET7a and SISWEET14 regulate sugar transport and storage in tomato fruits. *Horticulture Research* 8:186
 44. Yu M, Chen L, Liu D, Sun D, Shi G, et al. 2022. Enhancement of photosynthetic capacity in spongy mesophyll cells in white leaves of *Actinidia kolomikta*. *Frontiers in Plant Science* 13:856732
 45. Zhang X, Wen B, Zhang Y, Li Y, Yu C, Peng Z, et al. 2022. Transcriptomic and biochemical analysis reveal differential regulatory mechanisms of photosynthetic pigment and characteristic secondary metabolites between high amino acids green-leaf and albino tea cultivars. *Scientia Horticulturae* 295:110823



Copyright: © 2024 by the author(s). Published by Maximum Academic Press, Fayetteville, GA. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.