

Transcriptome and yeast two-hybrid sequencing shed light on the crosstalk between light and jasmonate signaling in regulating tanshinone biosynthesis

Han Zheng^{1,2#}, Xueqing Fu^{2#}, Muyao Yu^{1,3#}, Qi Liu^{1,4}, Chenyi Li⁵, Ling Li², Shuyi Qian², Kang Chen¹, Shuncang Zhang^{6*}, Kexuan Tang^{2*} and Luqi Huang^{1*}

¹ State Key Laboratory for Quality Ensurance and Sustainable Use of Dao-di Herbs, National Resource Center for Chinese Materia Medica, China Academy of Chinese Medical Sciences, Beijing 100700, China

² Frontiers Science Center for Transformative Molecules, Joint International Research Laboratory of Metabolic & Developmental Sciences, Plant Biotechnology Research Center, Fudan-SJTU-Nottingham Plant Biotechnology R & D Center, School of Agriculture and Biology, Shanghai Jiao Tong University, Shanghai 200240, China

³ School of Traditional Chinese Medicine, Beijing University of Chinese Medicine, Beijing 102401, China

⁴ College of Pharmacy, Nanjing University of Chinese Medicine, Nanjing 210023, China

⁵ Department of Biochemistry, Institute of Plant Biology, School of Life Sciences, Fudan University, Shanghai 200438, China

⁶ College of Bioscience and Biotechnology, Yangzhou University, Yangzhou 225009, China

Authors contributed equally: Han Zheng, Xueqing Fu, Muyao Yu

* Corresponding authors, E-mail: zhangsc@yzu.edu.cn; kxtang@sjtu.edu.cn; huanglq@cacms.cn

Abstract

As a high-value active component in *Salvia miltiorrhiza* (Danshen), the accumulation of tanshinones is under the influence of various environmental factors, yet it is unknown whether light, as an essential environmental factor for plant growth, regulates the biosynthesis of tanshinones. In this study, we first measured the temporal accumulation of jasmonates (JAs) and tanshinones under dark-to-light transition, and found that the accumulation of JA, JA-Ile and tanshinones is light-dependent. Transcriptome analysis revealed that *SmAOS* and *SmAOC* involved in JA biosynthesis responded rapidly under dark-to-light transition, followed by the accumulation of JA and JA-Ile. A light-responsive transcription factor *SmHY5* was filtered indicating a potential regulatory relationship with tanshinone-related genes. Co-expression analysis identified multiple classes of transcription factors highly correlated with JAs and tanshinone biosynthetic genes, among which *SmHY5* was confirmed to interact with *SmBBX21* through Y2H-seq. Further confirmation of the interaction between *SmHY5* and the IV subfamily of *SmBBXs*, including *SmBBX20-24*, was obtained through Y2H experiments. Specifically, *SmBBX20* and *SmBBX21* showed tissue-specific expression in aboveground and underground tissues, consistent with the accumulation sites of JAs and tanshinones. Taken together, this study contributes to the understanding of light and JA signaling in tanshinone biosynthesis in Danshen. We propose a crosstalk between light and JA signaling which regulates tanshinone biosynthesis through the *SmHY5-SmBBXs* complex and thus provide a basis for improving tanshinone production through the combination of light and JA.

Citation: Zheng H, Fu X, Yu M, Liu Q, Li C, et al. 2024. Transcriptome and yeast two-hybrid sequencing shed light on the crosstalk between light and jasmonate signaling in regulating tanshinone biosynthesis. *Medicinal Plant Biology* 3: e006 <https://doi.org/10.48130/mpb-0024-0006>

Introduction

Tanshinones are a group of abietane diterpenoid natural products synthesized in the root of *Salvia miltiorrhiza* Bunge, commonly called Danshen, a renowned traditional Chinese medicine. To date, over 40 tanshinones and structurally-related compounds have been identified from Danshen^[1], such as Cryptotanshinone (CPT), Tanshinone IIA (Tan IIA), Dihydrotanshinone I (DHT), and Tanshinone I (Tan I), which reportedly provide clinical benefits for treating cardiovascular diseases^[2]. The accumulation of tanshinones is frequently induced by the surrounding biotic and abiotic stresses, and is mediated by multi-tiered hormonal signaling pathways and transcription factors^[3]. Among these, jasmonates (JAs), including 12-oxo-phytodienoic acid (OPDA), jasmonic acid (JA), methyl jasmonate (MeJA), and jasmonoyl-isoleucine (JA-Ile), play crucial roles

in the regulation of tanshinone metabolism^[4]. JA-Ile, as a core small molecule in JA signaling transduction, plays a predominant role in connecting coronatine Insensitive 1 (COI1) with jasmonate-ZIM domain (JAZ) proteins. It promotes the ubiquitination and degradation of JAZ proteins, leading to the release of transcription factors and the regulation of downstream biosynthetic genes^[5]. For instance, *SmMYC2a/b* and *SmMYB39* interact with *SmJAZ1/2*, *SmERF73* interacts with *SmJAZ3*, *SmbHLH37* interacts with *SmJAZ3/8*, which ultimately regulate the biosynthesis of tanshinones and phenolic acids^[4,6–8]. Additionally, in *Artemisia annua*, the interaction between *AaWRKY9* and *AaJAZ9* has also been discovered^[9]. The types of transcription factors involved in JA signaling transduction through interactions with JAZ proteins are gradually being expanded.

As a vital environmental factor affecting plant growth and development, light is involved in various physiological

processes such as photomorphogenesis, lateral root development, seed germination and secondary metabolites accumulation^[10–12]. It was reported that white light treatment promotes content of DHT, red and blue light treatment significantly improves salvianolic acid content^[13,14]. Further investigation into the molecular mechanisms has revealed that light can indirectly promote the generation of the JA-Ile signaling molecule through the photoreceptors CRY1/2 and phyA/B^[10,15] and in turn, leads to the degradation of JAZ proteins and the subsequent release of MYCs transcription factors, which activate ELONGATED HYPOCOTYL 5 (HY5) transcription factor^[16]. HY5 further regulates downstream target genes to achieve the transmission of light signaling. These findings shed light on the complex relationship between light and JA signaling pathway in regulating plant growth under environmental stress.

HY5, a bZIP transcription factor, can be activated by photoreceptors to promote downstream photomorphogenesis. As a core regulatory factor regulating plant development, HY5 also participates in the biosynthesis of active compounds^[17]. It was reported that AtHY5 affects anthocyanin accumulation by interacting with AtPIF3^[18], regulates artemisinin biosynthesis by binding the ACGT motif on the *AaGSW1* promoter^[19] and meanwhile directly binds with G-box on the promoter of *AaQH6*, which encodes monoterpene pinene biosynthesis^[20]. BBX (B-box) transcription factor is a zinc finger protein extensively participating in plant life processes^[21], and is confirmed as a rate-limiting cofactor of HY5^[22]. BBX IV group PpBBX18 and PpBBX21 antagonistically regulates anthocyanin biosynthesis via competitive interaction with PpHY5^[23], OsBBX14 promotes photomorphogenesis by activating expression of OsHY5L1^[24], AtBBX11 and AtBBX21 regulates plant development by forming a positive feedback loop with AtHY5^[25]. Both HY5 and BBX have been reported to be associated with multiple signaling pathways in a dominant way, including the abscisic acid (ABA)^[26–28], gibberellin (GA)^[29,30], and brassinosteroid (BR)^[31] pathways. However, it remains unknown whether there is a crosstalk between light and JA signaling, which could potentially regulate the formation of the HY5-BBX complex.

In this study, the temporal changes in JAs and tanshinone content after dark-to-light transition were measured, confirming the positive regulatory effect of light on both components. Transcriptome analysis was conducted at corresponding time points to analyze the expression patterns of genes involved in JAs and tanshinone biosynthesis. A co-expression analysis network was then constructed based on the expression patterns of transcription factors and biosynthetic genes, along with the types of cis-regulatory elements in promoters, inferring potential regulatory relationships between SmHY5-SmBBXs and tanshinone. Using Y2H-seq technology, SmHY5 bait was employed to screen and identify SmBBX21 from a yeast library of Danshen, and strong interactions were also confirmed between SmBBX20/22/23/24 of BBX IV group and SmHY5, therefore suggesting the presence of a potential regulatory network from light-JA signaling to SmHY5-SmBBXs and finally to tanshinone in Danshen. However, further validation is required to elucidate the regulatory relationships between downstream target genes of SmHY5 and SmBBXs, which would ultimately reveal the molecular mechanisms of crosstalk regulation between light and JA signaling on tanshinone biosynthesis.

Results

Dark-to-light transition modulates accumulation of JAs and tanshinones

In order to investigate the effect of light signaling on the regulation of JAs and tanshinones in Danshen, we designed a light-induced experiment and treated 4-week-old plants with dark-to-light transition (Fig. 1a). The results showed that OPDA, as a substrate for JAs, exhibited a transient increase in both leaves and roots, reaching its peak at 4 h after treatment (Fig. 1b). In leaves, the content of OPDA increased dramatically to 98.7 ng/g, which was 3.0 times higher than that at 0 h (CK), while in roots, it increased significantly to 88.0 ng/g, which was 1.8 times higher than that at 0 h (CK), and then gradually decreased to a slightly higher level than that before treatment. JA was not detected in leaves at 0 h, but its content level gradually increased and reached its peak at 32 h with a gradient increase, reaching 270.3 ng/g; however, in roots, it reached its peak at 4 h, which was 3.4 times higher than that at 0 h (CK), with a content of 220.6 ng/g, and then gradually decreased. The signaling molecule of the JA pathway, JA-Ile, was not detected in leaves at 0 h, but its content level gently increased and reached its peak at 32 h, showing consistent trend with JA, with a content of 57.3 ng/g, while there was no obvious fluctuation in the content of JA-Ile in roots.

The content of diterpenoids also showed a trend of transient changes in response to dark-to-light transition (Fig. 1c). CPT was at a low level at 0 and 0.5 h, and increased abruptly at 1 h, then reached its peak at 4 h with a content of 0.47 mg/g, which was 4.7 times higher than that at 0 h (CK), and then gradually decreased but still significantly higher than that at 0 h (CK). There was a significant accumulation at 32 h. The content of Tan IIA also increased sharply at 1 h, reaching its peak at 4 h with a content of 0.54 mg/g, which was 2.5 times higher than that at 0 h (CK). DHT accumulated in a similar manner as CPT, increasing sharply at 1 h and reaching its peak at 4 h with a content of 0.029 mg/g, which was 4.5 times higher than that at 0 h (CK), and showed a significant accumulation at 32 h. Tan I accumulated in a similar way as Tan IIA, increasing sharply at 1 h and reached its peak at 4 h with a content of 0.065 mg/g, which was 1.9 times higher than that at 0 h (CK), and then gradually decreased. These results indicate that alteration in light can positively induce the elevation in content levels of JAs and diterpenoids, but the specific molecular mechanisms are still unclear.

Dark-to-light transition induces expression of JAs and tanshinone biosynthetic genes

To further investigate the effect of light on JA biosynthesis in Danshen, we measured the transcriptome after dark-to-light treatment (Fig. 2a). Among the six key genes involved in JA biosynthesis, the upstream *SmLOX* showed a decreasing trend in expression in both leaves and roots, while the two rate-limiting enzyme genes, *SmAOS* and *SmAOC*, were significantly upregulated in roots. *SmAOS* exhibited transient high expression in both leaves and roots within a short time, peaking at 0.5 h with an 8.7-fold and 1.8-fold increase, respectively. In leaves, *SmAOC* also reached its peak at 0.5 h with a 9.1-fold increase compared to the control, while in roots, it showed a continuous upward trend, ensuring the sustained supply of JA precursor OPDA. The expression of *SmOPR3* was relatively stable in




Fig. 1 Transient increase in the accumulation of JAs and tanshinones after dark-to-light transition in Danshen. (a) The schematic representation depicts the dark-to-light transition treatment. (b) Contents of 12-oxo-phytodienoic acid (OPDA), jasmonic acid (JA), jasmonic acid-isoleucine (JA-Ile) in root and leaf determined by UPLC-MS/MS. (c) Contents of Cryptotanshinone I (CPT), Tanshinone IIA (Tan IIA), Dihydrotanshinone (DHT), and Tanshinone I (Tan I) in root determined by UPLC-MS. Data are shown as the mean \pm SD ($n = 6$). Variance test was performed for multigroup comparison, with different lowercase letters indicating significant difference between groups ($p < 0.05$).

both leaves and roots, with a slight increase in roots (Supplemental Table S1). Among the downstream signaling molecule biosynthesis genes, the peak expression level of *SmJAR1* was later than that of *SmAOS* and *SmAOC*, reaching its highest level at 2 h in leaves, which was 2.2 times higher than that of the control, but remained relatively stable in roots, possibly due to the stable content of JA-Ile in roots. In leaves, the expression level of *SmJMT* also peaked at 2 h, reaching 5.1 times that of the control, but was not detected in roots. More importantly, the light-responsive transcription factor *HY5* was activated and expressed within a short period of time, reaching its peak at 1 h in leaves, which was 76.1 times higher than the control. Therefore, light promoted the expression of most JA biosynthesis genes in Danshen, especially *SmAOS* and *SmAOC*, but the causal relationship and the complex synergistic regulation mechanism on tanshinone accumulation still need further exploration.

To investigate the effect of light on tanshinone biosynthesis, we evaluated the expression levels of key enzyme genes in the biosynthetic pathways (Fig. 2b). Regarding the genes involved in obtaining IPP and DMAPP via the MVA and MEP pathways, there was an overall increasing trend. However, in the MVA pathway, the rate-limiting enzyme gene *SmHMGS* reached its highest expression level in both leaves and roots at 1 h, while another rate-limiting enzyme gene *SmHMGR* showed a continuously decreased expression in both tissues. This opposite expression trend may limit the supply of precursors for diterpene biosynthesis through the MVA pathway. Nevertheless, the genes *SmMK*, *SmPMK*, and *SmMVD* still exhibited distinct expression peaks within a short time in both leaves and roots.

In the MEP pathway, the expression trends of the rate-limiting enzyme genes *SmDXS2* and *SmDXR1* were more consistent, reaching their highest levels at 1 and 2 h, respectively, ensuring the supply of precursors for subsequent diterpenoid compounds. Subsequently, the genes *SmMCT*, *SmCMK*, *SmMDS*, *SmHDS*, *SmHDR*, and *SmIDI* also showed varied degrees of upregulation. Of particular importance, during the formation of GGPP, the substrate for tanshinone biosynthesis, the gene *GGPPS* responded positively to light signaling in both leaves and roots, reaching its peak at 0.5 and 2 h, respectively. Subsequently, the genes involved in diterpene synthase and *SmCYP450* enzymes exhibited specific high expression in roots, with *SmCPS1*, *SmKSL1*, and *SmCYP76AH3* reaching their highest levels at 1 h. The expression peaks of *CYP450* genes involved in late-stage modifications generally occurred later, with *SmCYP76AK1*, *SmCYP71D373*, and *SmTIAS* gradually increasing and peaking at 32 h. These changes in gene expression ultimately reflected in the variation of tanshinone content. However, the molecular mechanism by which light signaling regulates tanshinone biosynthesis is still unclear and requires further research.

Multiple transcription factors potentially regulate the biosynthesis of JAs and tanshinones in the process of dark-to-light transition

As transcription factors play essential roles in plant secondary metabolite accumulation and growth, the upstream regulatory network of JAs and tanshinone accumulation was analyzed. Co-expression analysis revealed significant co-expression between genes involved in JAs (*SmLOX*, *SmAOS*, *SmAOC*, *SmOPR3*, *SmJAR*, *SmJMT*) and transcription factors such as




Fig. 2 Dark-to-light transition affects biosynthesis of both (a) JAs and (b) tanshinones. Heat map showing expression fold changes (\log_2 FC) of genes and FPKM values are shown in [Supplemental Table S1](#). (a) α -LNA, α -Linolenic Acid; 13-HPOT, 13(s)-hydroperoxy-linolenic acid; 12,13-EOT, 12,13(S)-epoxy-9(Z),11,15(Z)-octadecatrienoic acid; OPDA, 12-oxo-phytyldienoic acid; OPC-8:0, 3-oxo-2-(2-(Z)-pentenyl)-cyclopentane-1-octanoic acid; JA, jasmonic acid; JA-Ile, jasmonic acid-isoleucine; MeJA, methyl jasmonate; 13-LOX, 13-Lipoxygenase; AOS, allen oxidase; AOC, allene oxide cyclase; OPR, 12-oxophytyldienoate reductase; JAR, JA-amino acid synthetase; JMT, jasmonate hydroxymethyl transferase; HY5, elongated hypocotyl 5. (b) MVA pathway, the mevalonate pathway; Acetyl-CoA, acetyl coenzyme A; Acetoacetyl-CoA, acetoacetyl coenzyme A; HMG-CoA, 3-hydroxy-3-methylglutaryl CoA; MVA, mevalonate; MVA-P, mevalonate-5-phosphate; MVA-PP, mevalonate-5-diphosphomevalonate; IPP, isopentenyl pyrophosphate; MEP pathway, the 2-C-methyl-d-erythritol 4-phosphate pathway; G3P, glyceraldehyde 3-phosphate; DXP, 1-deoxy-d-xylulose 5-phosphate; MEP, 2-C-methyl-d-erythritol 4-phosphate; CDP-ME, 4-(cytidine 5'-diphospho)-2-C-methyl-d-erythritol; PP-CDP-ME, 2-phospho-4-(cytidine 5'-diphospho)-2-C-methyl-d-erythritol; ME-cPP, 2-C-methyl-d-erythritol 2,4-cyclodiphosphate; H-ME-B-PP, 1-hydroxy-2-methyl-2-butenyl 4-diphosphate; DMAPP, dimethylallyl pyrophosphate; GGPP, geranylgeranyl diphosphate; normal-CPP, normal-copalyl diphosphate; AACT, acetoacetyl-CoA transferase; HMGS, hydroxymethyl-glutaryl-CoA synthase; HMGR, 3-hydroxy-3-methylglutaryl CoA reductase; MK, mevalonate kinase; PMK, phosphomevalonate kinase; MVD, mevalonate-5-pyrophosphate decarboxylase; DXS, 1-deoxy-d-xylulose-5-phosphate synthase; DXR, 1-deoxy-d-xylulose-5-phosphate reductoisomerase; MCT, 2-C-methyl-D-erythritol 4-phosphate cytidyltransferase; CMK, 4-(cytidine 5'-diphospho)-2-C-methyl-d-erythritol kinase; MDS, 2-C-methyl-d-erythritol 2,4-cyclodiphosphate synthase; HDS, 4-hydroxy-3-methylbut-2-enyl diphosphate synthase; HDR, 4-hydroxy-3-methyl-but-2-enyl diphosphate reductase; IDI, Isopentenyl-diphosphate delta-isomerase; GGPPS, GGPP synthase; CPS1, copalyl diphosphate synthase; KSL1, kaurene synthase-like; TIIAS, tanshinone IIA synthase.

WRKYs, *ERFs*, *MYBs*, *bHLHs*, *bZIPs*, *BBXs* (Fig. 3a). Similarly, genes related to tanshinone biosynthesis (*SmDXS2*, *SmDXR1*, *SmGGPPS*, *SmCPS1*, *SmKSL1*, *SmCYP76AH1*, *SmCYP76AH3*, *SmCYP76AK1*, *SmCYP71D375*, *SmCYP71D373*, *SmTIIAS*) exhibited co-expression with these transcription factors (Fig. 3b).

Further analysis revealed a high correlation ($\text{Cor} > 0.75$) between *SmHY5* and key genes in the MVA pathway, such as *SmHMGS*, *SmMK*, as well as genes in the MEP pathway, including *SmDXR1* and *SmIDI* (Fig. 3c). Furthermore, *SmHY5* showed a

moderate correlation ($\text{Cor} > 0.5$) with terpene synthase genes like *SmCPS1* and *SmKSL1*, as well as CYP450 genes like *SmCYP76AH1* and *SmCYP71D373*. However, no correlation was found between *SmHY5* and genes involved in JA biosynthesis, suggesting that *SmHY5* is more likely to regulate upstream terpenoid precursor forming related genes.

On the other hand, *SmBBX20* exhibited higher expression level in leaves and showed strong correlations ($\text{Cor} > 0.75$) with JA biosynthesis genes such as *SmAOS* and *SmAOC* (Fig. 3c). The




Fig. 3 JAs and tanshinone biosynthetic genes potentially regulated by transcription factors. (a) Co-expression analysis of JAs biosynthetic genes and transcription factors in leaf. (b) Co-expression analysis of tanshinone biosynthetic genes and transcription factors in root, while pathway genes are represented by octagons and transcription factors by circles. The screening criterion for selection was set as a correlation coefficient greater than 0.3. (c) Transcription factor-pathway genes with correlation coefficients, green represents genes in leaves and yellow in roots. (d) Cis-elements of the JAs biosynthetic gene promoters. (e) Cis-elements of the tanshinone biosynthetic gene promoters.

expression levels of these genes were consistent with BBX20 and all peaked at 0.5 h (Supplemental Table S1). *SmBBX20* also displayed moderate correlations ($\text{Cor} > 0.5$) with MEP pathway genes like *SmPMK*, *SmHDR*, as well as diterpene synthase genes *SmGGPPS*, *SmKSL1*, and *SmCYP76AH1*, *SmCYP76AH3*. In roots, it showed a high correlation ($\text{Cor} = 0.81$) with MEP pathway gene *SmMCT*, and moderate correlations ($\text{Cor} > 0.5$) with *SmCYP76AK1* and *SmTIAS*. Interestingly, *SmBBX20* also demonstrated a strong correlation with *SmBBX21*, indicating a possible regulatory relationship. In leaves, *SmBBX21* displayed high correlations ($\text{Cor} > 0.75$) with MVA pathway genes like *SmMK*, *SmIDI*, *SmMVD*, and *SmHMGs*, as well as moderate correlations ($\text{Cor} > 0.5$) with MEP pathway genes like *SmMDS2*, *SmHDR2*, and diterpene synthase genes *SmCPS1*, *SmKSL1*, as well as *SmCYP76AH1*, *SmCYP71D375*. Notably, *SmBBX21* also showed

moderate correlations ($\text{Cor} > 0.5$) with JA biosynthesis genes like *SmAOS* and *SmAOC*, as well as *SmHY5*. However, *SmBBX21* exhibited lower expression levels in leaves compared to roots, where it displayed strong correlations ($\text{Cor} > 0.75$) with *SmCYP71D373*, *SmTIAS*, and moderate correlations ($\text{Cor} > 0.5$) with MEP pathway genes like *SmMCT*, *SmHDR1/2*. Additionally, the highest correlation of 0.91 was observed between *SmBBX21* and *SmJAR*. Therefore, it is proposed that *SmBBX20* and *SmBBX21* potentially possess the ability to cooperatively regulate the biosynthesis of JAs and tanshinones.

Subsequently, based on the published genomic data, we cloned and analyzed the cis-elements in the promoter regions of JAs and tanshinone biosynthetic genes^[32,33]. As shown in Fig. 3d & e, the promoter regions of genes involved in JAs and tanshinone biosynthesis contain multiple cis-elements that are

binding sites for the above-mentioned transcription factors. These cis-elements include the G-box for HY5 (bZIP), T/G-box for BBX, E-box for bHLH, W-box for WRKY, MBS for MYB, and GCC-box for ERF, providing possibilities for direct or indirect signal transduction and transcriptional regulation.

SmHY5 specifically interacts with SmBBXs in the roots of Danshen

In order to uncover the proteins that interact with SmHY5, we built SmHY5 into a construct using pGBKT7 as back bone to screen Y2H library of Danshen root tissue. After ensuring no self-activation in pGBKT7-SmHY5, we collected 192 positive clones on SD-Trp/Leu/His on which we performed PCR single-cloning sequencing. Through BLAST analysis to eliminate duplicate sequences, we obtained 63 target genes (Supplemental Table S2), among which were several transcription factors, including SmWRKY1^[34], SmWRKY41, SmMYB1R1, SmbHLH3^[35], and SmBBX21. Further validation revealed that 13 clones, including SmbHLH3 and SmBBX21, were able to grow on SD-Trp/Leu/His/Ade plates and displayed blue color on SD-Trp/Leu/His/Ade + X- α -gal plates (Fig. 4a). Based on SmBBX21's classification in the IV subfamily of BBX, we proceeded to screen for potential interactions between proteins in this

subfamily and SmHY5. Seven genes from this subfamily, *SmBBX18*, *SmBBX19*, *SmBBX20*, *SmBBX21*, *SmBBX22*, *SmBBX23*, *SmBBX24*, were cloned and then built into the constructs for yeast two-hybrid. From the results of -Trp/Leu and -Trp/Leu/His/Ade culture medium, the physical interactions between SmHY5 and SmBBX20, SmBBX21, SmBBX22, SmBBX23, SmBBX24 were proved, which can connect light pathway and the BBX gene family (Fig. 4b).

In order to validate the potential role of SmBBXs in regulating tanshinone biosynthesis in root, we conducted tissue-specific expression analysis on Danshen. We found that *SmBBX20* exhibits higher expression levels in leaves, stems, and flowers, while *SmBBX21* shows specific expression in root tissues, including rhizome, fibrous roots, cortex, and xylem (Fig. 4c). This may be associated with the regulatory role of SmBBXs in different tissues.

Discussion

Tanshinones, a class of valuable active components found in Danshen, are biosynthesized under the induction of various biotic and abiotic stresses. JAs serve as important hormone signaling molecules that respond to stress and act as 'brokers'




Fig. 4 Potential interaction between root-specific expressed SmBBX21 and SmHY5. (a) Yeast two-hybrid sequencing (Y2H-seq) screened potential interacting proteins of SmHY5 in root. Monoclonal sequencing results are shown in Supplemental Table S2. (b) Y2H confirmed the interaction between SmHY5 and SmBBXs. (c) The expression of the seven candidate *SmBBX* genes in different tissues of Danshen. *ACTIN* was used as an internal control; for each gene, the expression level relative to *ACTIN* in immature seed was taken as 1. Data are shown as the mean \pm SD ($n = 3$).

Light-jasmonate crosstalk regulates tanshinones

linking the environment to secondary metabolism^[3]. Among them, JA-Ile can promote the degradation of JAZ, release transcription factors, and activate downstream biosynthetic genes^[4,6–9].

This study found that the important signaling molecule JA-Ile was not detected in leaves under dark conditions, but increased steadily after exposure to light, indicating that JA-Ile biosynthesis is light-dependent (Fig. 1b). Studies have reported that light can indirectly promote the generation of JA-Ile signaling molecules through photoreceptors CRY1/2 and phyA/B,

and then regulate HY5 through MYCs transcription factors^[10,15,16]. In this study, the two upstream key enzyme genes *SmAOS* and *SmAOC* in Danshen JAs biosynthetic pathway reached their peak at 0.5 h, and JA-Ile was detected and accumulated within the same time frame (Fig. 2a). This time point was prior to the high expression time point of *SmHY5*, which also suggests the possibility of indirect regulation of *SmHY5* by JAs in Danshen.

Furthermore, HY5 can directly bind to the promoters of light-responsive transcription factors and biosynthetic genes,




Fig. 5 Hypothesis for crosstalk between light and JA signaling regulating tanshinone biosynthesis through the SmHY5-SmBBXs complex. Under dark conditions, the absence of JA-Ile leads to the inhibition of multiple transcription factors by SmJAZ proteins. Simultaneously, SmCOP1 degrades SmHY5, preventing the formation of a regulatory complex between SmHY5 and SmBBXs. Consequently, the expression of downstream genes involved in tanshinone biosynthesis is at a lower level. In contrast, under light conditions, JA-Ile biosynthesis is enabled, leading to the degradation of SmJAZ proteins and the release of SmMYC and other transcription factors. Additionally, SmCOP1 releases SmHY5, allowing the formation of a regulatory complex between SmHY5 and SmBBXs. This coordination activates the expression of downstream genes involved in tanshinone biosynthesis. Finally, the crosstalk between light and JA signaling on tanshinone biosynthesis was realized through the formation of SmHY5 and SmBBXs complex or not.

thereby regulating plant metabolism^[17]. For example, AaHY5 activates the transcription of *AaWRKY9/14*, promoting artemisinin accumulation^[9,36]. SmHY5 directly binds to *SmC4H1* in hairy roots of Danshen, enhancing phenolic acid accumulation^[37]. In this study, co-expression analysis revealed a strong correlation ($\text{cor} > 0.5$) between SmHY5 and various transcription factors, as well as tanshinone biosynthetic genes (Fig. 3c). Multiple G-box elements were identified in the promoters of these genes, suggesting their potential regulation by SmHY5 (Fig. 3e). However, it has been reported that the direct binding ability of HY5 to gene promoters is weak and requires the formation of heterodimers with BBX proteins for co-regulation with HY5-COP1 cascade module^[38]. Among them, BBXs are believed to interact directly with HY5, exerting promoting or inhibitory effects on HY5 function^[21]. The relationship between BBX IV group and HY5 is particularly close^[39], with not only a direct interaction between AtBBX20 and AtHY5 but also a mutual effect on their protein abundance^[40]. AtBBX21 interacts with AtHY5 and meanwhile exerts transcriptional regulation on AtHY5^[41]. In this study, Y2H-seq was utilized to screen and identify SmBBX21 from a yeast library of Danshen roots using SmHY5 as bait (Fig. 4a). Furthermore, strong interactions were also confirmed between SmHY5 and members of BBX IV group, namely SmBBX20-24 (Fig. 4b). SmBBX20 showed a peak correlation with *SmAOS*, *SmAOC*, and *SmJAR* at 0.5 h (Fig. 2a, 3c), and it was specifically expressed in the aerial parts (Fig. 4c), suggesting its potential involvement in JA biosynthesis regulation. The peak of SmBBX21 expression occurred after 2 h (Supplemental Table S1), showing specific expression in the underground parts (Fig. 4c) and exhibiting strong correlations with MEP pathway and tanshinone biosynthetic genes (Fig. 3c). Based on the promoter G-box and T/G-box elements, it is speculated that there may be regulation of tanshinone biosynthesis by SmBBX21.

Therefore, a potential regulatory network from light-JA signaling to SmHY5-SmBBXs to tanshinone biosynthesis is constituted (Fig. 5). However, further experimental validation is needed for the downstream target genes of SmHY5 and SmBBXs, as well as the regulatory effects on tanshinone biosynthetic genes. Subsequent research is required to uncover the molecular mechanisms underlying the crosstalk regulation of tanshinone by the light and JA signaling. This study supplements the information on light and JA signaling in Danshen as a medicinal model plant and provides a basis for improving tanshinone production through the combination of light and JA.

Conclusions

Our research reports the positive regulatory effect of light on the accumulation of JAs and tanshinones, and analyzes the changing patterns of expression levels of genes involved in JAs and tanshinone biosynthesis after the transition from darkness to light. We identified transcription factors highly correlated with the aforementioned biosynthetic genes, including HY5 and BBXs. The interactions between HY5 and BBX IV group BBX20-24 were confirmed through Y2H-Seq and Y2H, and the tissue-specific expression of BBXs was also observed. Thus, we proposed a potential regulatory network in which the crosstalk between light and JA signaling regulates tanshinone biosynthesis through the SmHY5-SmBBXs complex.

Materials and methods

Plant materials

The plants of Danshen were cultivated and stored in the greenhouse of National Resource Center for Chinese Materia Medica, China Academy of Chinese Medical Sciences (Beijing, China). The growth conditions were maintained at 25 °C with a 16-h light period each day for four weeks, followed by 48 h of darkness before the light treatment.

RNA-seq and analysis

Total RNAs treated with light (0–36 h) were extracted as described earlier^[4]. RNA-seq library preparation and paired-end sequencing were both performed using an Illumina HiSeq 6000 platform (Novogene, Beijing, China). FeatureCounts was adopted to quantify total expressed transcripts with the FPKM (fragments per kilobase of exon model per million mapped reads) value^[42]. Differentially expressed genes (DEGs) were obtained by comparing FPKM value among 0–36 h samples. DEGs with a \log_2 fold-change ($|\log_2 \text{FC}|$) ≥ 1 and an adjusted q -value ≤ 0.05 were identified using the DEG_{SEQ} R package^[43]. GENIE3 R package was used to calculate the Pearson correlation coefficients between transcription factors^[44] and genes in batches and Cytoscape software (version 2.6.3) to draft the correlation map.

Phytohormone determination

Phytohormones including JA, OPDA, JA-Ile were quantitated using a previously reported method^[45]. Fifty mg of ground plant material was subjected to extraction using an isopropanol-water-hydrochloric acid solution (2:1:0.002, V/V/V) for a duration of 30 min. Subsequently, the extract was further extracted with dichloromethane for another 30 min. The organic layer obtained was evaporated under a stream of nitrogen until dryness, resulting in a residue. This residue was then dissolved in 70% methanol to obtain a solution. To remove any insoluble particles, the solution was subjected to centrifugation at 12,000 rpm for 15 min, resulting in the collection of the supernatant. The quantitation of endogenous phytohormone was conducted using an ACQUITY UPLC I-Class system equipped with a 6500 QTRAP mass spectrometer. The mobile phase consisted of acetonitrile (A) and 0.05% formic acid-water (B), and the flow rate was set at 0.4 mL/min. The gradient elution was as follows: 0–3.5 min, 10% A; 3.5–4 min, 10%–95% A; 4–8 min, 95% A. For analysis, 1 μL of the sample solution was injected into the system. Mass spectrometry analysis was performed in both positive and negative modes using multiple reaction monitoring (MRM) mode.

Tanshinones determination

Leaves and roots were collected and lyophilized overnight using an FDU-1110 freezer dryer (Eyela, Tokyo, Japan). The samples were accurately weighed and sonicated in methanol at a ratio of 1:30 (w/v) for 30 min using an SB-800 DTD sonicator (Ningbo Xinzhi, China; power, 100 W; frequency, 40 kHz). The resulting supernatant was filtered through a 0.22 μm filter (Pall, Ann Arbor, MI, USA) for subsequent ultra-performance liquid chromatography coupled with triple-quadrupole tandem mass spectrometry (UPLC-TQ-MS) analysis. Experiments included six biological replicates and the statistical significance was evaluated using a variance analysis in SPSS v.16.0^[46].

Quantitative real-time PCR analysis

To evaluate the expression profiles of the *SmBBX* genes in different tissues, a qPCR assay was conducted in a 10 μ l reaction volume containing 5 μ l of TB Green® Premix Ex Taq (TaKaRa RR820B, Japan), 0.5 μ M of each oligonucleotide and 1 μ l of cDNA template on a StepOnePlus PCR system (Roche LightCycler 480). The specific qPCR primers used in this study are listed in [Supplemental Table S3](#). The β -actin genes were utilized as endogenous controls. Three technical replicates were performed for each sample, and quantitative analysis was performed using the $2^{-\Delta\Delta CT}$ method^[47].

Yeast two-hybrid assays

The yeast two-hybrid screening method was performed according to the manufacturer's user guide (Takara Bio Inc.). The full-length cDNA of SmHY5 was amplified and cloned into the pGBKT7 vector, which was then transformed into the yeast strain AH109. This construct served as the bait to screen a cDNA library prepared from equal amounts of poly(A)-containing RNA extracted from root tissue. The screening and plasmid extraction sequencing steps were performed following the manufacturer's instructions. The full-length cDNAs of *SmBBXs* were amplified and subcloned into the pGADT7 vector individually. The pGADT7 and pGBKT7 constructs were co-transformed into the yeast strain AH109, and the transformed cells were selected on SD/-Leu/-Trp medium. The resulting co-transformed cells were streaked on quadruple dropout medium (SD/-His/-Leu/-Trp/-Ade) and incubated at 30 °C for 72 h.

Author contributions

The authors confirm contribution to the paper as follows: study conception and design: Huang L, Zheng H; analysis and interpretation of results: Zheng H, Fu X, Yu M, Liu Q, Li C, Li L, Qian S, Chen K; draft manuscript preparation: Zheng H, Fu X, Yu M; manuscript editing: Zhang S, Tang K; project supervision: Huang L. All authors reviewed the results and approved the final version of the manuscript.

Data availability

RNA-seq data have been deposited in the NCBI repository: SRR27169169- SRR27169184. Other data supporting the findings of this study are available within the paper and within its supplementary materials published online.

Acknowledgments

This research was funded by Natural Science Foundation of China (82104342), CACMS Innovation Fund (CI2021A041003, CI2023E002), National Key Research and Development Program of China (2023YFC3503900), Fundamental Research Funds for the Central public welfare research institutes (ZZ15-YQ-060), China Postdoctoral Science Foundation (2022M712099), key project at central government level: The ability establishment of sustainable use for valuable Chinese medicine resources (2060302).

Conflict of interest

The authors declare that they have no conflict of interest.

Supplementary information accompanies this paper at (<https://www.maxapress.com/article/doi/10.48130/mpb-0024-0006>)

Dates

Received 31 December 2023; Revised 17 February 2024; Accepted 26 February 2024; Published online 3 April 2024

References

- Mei XD, Cao YF, Che YY, Li J, Shang ZP, et al. 2019. Danshen: a phytochemical and pharmacological overview. *Chinese Journal of Natural Medicines* 17:59–80
- Jiang Z, Gao W, Huang L. 2019. Tanshinones, Critical Pharmacological Components in *Salvia miltiorrhiza*. *Frontiers in Pharmacology* 10:202
- Zheng H, Fu X, Shao J, Tang Y, Yu M, et al. 2023. Transcriptional regulatory network of high-value active ingredients in medicinal plants. *Trends in Plant Science* 28(4):429–46
- Zheng H, Jing L, Jiang X, Pu C, Zhao S, et al. 2021. The ERF-VII transcription factor SmERF73 coordinately regulates tanshinone biosynthesis in response to stress elicitors in *Salvia miltiorrhiza*. *New Phytologist* 231:1940–55
- Howe GA, Major IT, Koo AJ. 2018. Modularity in jasmonate signaling for multistress resilience. *Annual Review of Plant Biology* 69:387–415
- Zhou Y, Sun W, Chen J, Tan H, Xiao Y, et al. 2016. *SmMYC2a* and *SmMYC2b* played similar but irreplaceable roles in regulating the biosynthesis of tanshinones and phenolic acids in *Salvia miltiorrhiza*. *Scientific Reports* 6:22852
- Du T, Niu J, Su J, Li S, Guo X, et al. 2018. SmbHLH37 functions antagonistically with SmMYC2 in regulating jasmonate-mediated biosynthesis of phenolic acids in *Salvia miltiorrhiza*. *Frontiers in Plant Science* 9:1720
- Ma P, Pei T, Lv B, Wang M, Dong J, et al. 2022. Functional pleiotropism, diversity, and redundancy of *Salvia miltiorrhiza* Bunge JAZ family proteins in jasmonate-induced tanshinone and phenolic acid biosynthesis. *Horticulture Research* 9:uhac166
- Fu X, Peng B, Hassani D, Xie L, Liu H, et al. 2021. AaWRKY9 contributes to light- and jasmonate-mediated to regulate the biosynthesis of artemisinin in *Artemisia annua*. *New Phytologist* 231:1858–74
- Yi R, Yan J, Xie D. 2020. Light promotes jasmonate biosynthesis to regulate photomorphogenesis in *Arabidopsis*. *Science China Life Sciences* 63:943–52
- Alallaq S, Ranjan A, Brunoni F, Novák O, Lakehal A, et al. 2020. Light controls de novo adventitious root regeneration by modulating jasmonate and cytokinin homeostasis in Norway spruce hypocotyls. *bioRxiv* Preprint
- Wang Y, Fan X, Lin F, He G, Terzaghi W, et al. 2014. Arabidopsis noncoding RNA mediates control of photomorphogenesis by red light. *Proceedings of the National Academy of Sciences of the United States of America* 111:10359–64
- Feng S, Wang R, Gu W, Yu B, Wang Y, et al. 2019. Effects of red light and blue light on root morphology and accumulation of bioactive compounds in *Salvia miltiorrhiza*. *Chinese Traditional and Herbal Drugs* 50:5313–18
- Liang Z, Li Q, Xu W. 2012. Effects of different light quality on growth, active ingredients and enzyme activities of *Salvia miltiorrhiza*. *China Journal of Chinese Materia Medica* 37:2055–60
- Svyatyna K, Jikumaru Y, Brendel R, Reichelt M, Mithöfer A, et al. 2013. Light induces jasmonate - isoleucine conjugation via OsJAR1 - dependent and - independent pathways in rice. *Plant, Cell & Environment* 37:827–39

16. Ortigosa A, Fonseca S, Franco - Zorrilla JM, Fernández - Calvo P, Zander M, et al. 2020. The JA - pathway MYC transcription factors regulate photomorphogenic responses by targeting HY5 gene expression. *The Plant Journal* 102:138–52
17. Gangappa SN, Botto JF. 2016. The Multifaceted Roles of HY5 in Plant Growth and Development. *Molecular Plant* 9:1353–65
18. Shin J, Park E, Choi G. 2007. PIF3 regulates anthocyanin biosynthesis in an HY5-dependent manner with both factors directly binding anthocyanin biosynthetic gene promoters in Arabidopsis. *The Plant Journal* 49:981–94
19. Hao X, Zhong Y, Nützmann HW, Fu X, Yan T, et al. 2019. Light-Induced artemisinin biosynthesis is regulated by the bZIP transcription factor AaHY5 in *Artemisia annual*. *Plant and Cell Physiology* 60:1747–60
20. Zhou F, Sun T, Zhao L, Pan X, Lu S. 2015. The bZIP transcription factor HY5 interacts with the promoter of the monoterpene synthase gene QH6 in modulating its rhythmic expression. *Frontiers in plant science* 6:304
21. Gangappa SN, Botto JF. 2014. The BBX family of plant transcription factors. *Trends in Plant Science* 19:460–70
22. Bursch K, Toledo-Ortiz G, Pireyre M, Lohr M, Braatz C, et al. 2020. Identification of BBX proteins as rate-limiting cofactors of HY5. *Nature Plants* 6:921–28
23. Bai S, Tao R, Yin L, Ni J, Yang Q, et al. 2019. Two B-box proteins, PpBBX18 and PpBBX21, antagonistically regulate anthocyanin biosynthesis via competitive association with *Pyrus pyrifolia* ELONGATED HYPOCOTYL 5 in the peel of pear fruit. *The Plant Journal* 100:1208–23
24. Bai B, Lu N, Li Y, Guo S, Yin H, et al. 2019. OsBBX14 promotes photomorphogenesis in rice by activating OsHY5L1 expression under blue light conditions. *Plant Science* 284:192–202
25. Zhao X, Heng Y, Wang X, Deng X, Xu D. 2020. A positive feedback loop of BBX11-BBX21-HY5 promotes photomorphogenic development in *Arabidopsis*. *Plant Communications* 1:100045
26. Xu D, Gao S, Ma Y, Wang X, Feng L, et al. 2017. The G-Protein β subunit AGB1 promotes hypocotyl elongation through inhibiting transcription activation function of BBX21 in *Arabidopsis*. *Molecular Plant* 10:1206–23
27. Xu D, Li J, Gangappa SN, Hettiarachchi C, Lin F, et al. 2014. Convergence of light and ABA signaling on the ABI5 promoter. *PLoS Genetics* 10:e1004197
28. Chen H, Zhang J, Neff MM, Hong S, Zhang H, et al. 2008. Integration of light and abscisic acid signaling during seed germination and early seedling development. *Proceedings of the National Academy of Sciences of the United States of America* 105:4495–500
29. Wang F, Zhang L, Chen X, Wu X, Xiang X, et al. 2018. SIHY5 integrates temperature, light and hormone signaling to balance plant growth and cold tolerance. *Plant Physiology* 179:749–60
30. Weller JL, Hecht V, Vander Schoor JK, Davidson SE, Ross JJ. 2009. Light regulation of gibberellin biosynthesis in Pea is mediated through the COP1/HY5 pathway. *The Plant Cell* 21:800–13
31. Fan XY, Sun Y, Cao DM, Bai MY, Luo XM, et al. 2012. BZS1, a B-box protein, promotes photomorphogenesis downstream of both brassinosteroid and light signaling pathways. *Molecular Plant* 5:591–600
32. Ma Y, Cui G, Chen T, Ma X, Wang R, et al. 2021. Expansion within the CYP71D subfamily drives the heterocyclization of tanshinones synthesis in *Salvia miltiorrhiza*. *Nature Communications* 12:685
33. Li CY, Yang L, Liu Y, Xu ZG, Gao J, et al. 2022. The sage genome provides insight into the evolutionary dynamics of diterpene biosynthesis gene cluster in plants. *Cell Reports* 40:111236
34. Cao W, Wang Y, Shi M, Hao X, Zhao W, et al. 2018. Transcription factor SmWRKY1 positively promotes the biosynthesis of tanshinones in *Salvia miltiorrhiza*. *Frontiers in Plant Science* 9:554
35. Zhang C, Xing B, Yang D, Ren M, Guo H, et al. 2020. SmbHLH3 acts as a transcription repressor for both phenolic acids and tanshinone biosynthesis in *Salvia miltiorrhiza* hairy roots. *Phytochemistry* 169:112183
36. Zhou L, Huang Y, Wang Q, Guo D. 2021. AaHY5 ChIP-seq based on transient expression system reveals the role of AaWRKY14 in artemisinin biosynthetic gene regulation. *Plant Physiology and Biochemistry* 168:321–28
37. Deng C, Shi M, Fu R, Zhang Y, Wang Q, et al. 2020. ABA-responsive transcription factor bZIP1 is involved in modulating biosynthesis of phenolic acids and tanshinones in *Salvia miltiorrhiza*. *Journal of Experimental Botany* 71:5948–62
38. Yang G, Zhang C, Dong H, Liu X, Guo H, et al. 2022. Activation and negative feedback regulation of SIHY5 transcription by the SIBBX20/21–SIHY5 transcription factor module in UV-B signaling. *The Plant Cell* 34:2038–55
39. Li Y, Tong Y, Ye J, Zhang C, Li B, et al. 2023. Genome-Wide characterization of B-Box gene family in *Salvia miltiorrhiza*. *International Journal of Molecular Sciences* 24:2146
40. Wei C, Chien C, Ai L, Zhao J, Zhang Z, et al. 2016. The *Arabidopsis* B-box protein BZS1/BBX20 interacts with HY5 and mediates strigolactone regulation of photomorphogenesis. *Journal of Genetics and Genomics* 43:555–63
41. Xu D, Jiang Y, Li J, Lin F, Holm M, et al. 2016. BBX21, an *Arabidopsis* B-box protein, directly activates HY5 and is targeted by COP1 for 26S proteasome-mediated degradation. *Proceedings of the National Academy of Sciences of the United States of America* 113:7655–60
42. Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30:923–30
43. Wang L, Feng Z, Wang X, Wang X, Zhang X. 2010. DEGseq: an R package for identifying differentially expressed genes from RNA-seq data. *Bioinformatics* 26:136–38
44. Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. 2010. Inferring regulatory networks from expression data using tree-based methods. *PLoS One* 5:e12776
45. Chen K, Liu J, Ji R, Chen T, Zhou X, et al. 2019. Biogenic synthesis and spatial distribution of endogenous phytohormones and ginsenosides provide insights on their intrinsic relevance in *Panax ginseng*. *Frontiers in Plant Science* 9:1951
46. Frey F. 2017. SPSS (Software). In *The International Encyclopedia of Communication Research Methods*, eds. Matthes J, Davis CS, Potter RF. Los Angeles, USA: John Wiley & Sons. pp. 1–2. <https://doi.org/10.1002/9781118901731.iecrm0237>
47. Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* 25:402–08



Copyright: © 2024 by the author(s). Published by Maximum Academic Press, Fayetteville, GA. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.