

Chlorophyll-binding protein *AtSep2* overexpression causes leaf yellowing and enhances tolerance to high light stress in poplar

Donghui Chao¹, Xiaou Zhai², Lulu Ren¹, Shan Fu¹, Jin Zhang¹, Shuang Feng³, Haizhen Zhang^{1*} and Aimin Zhou^{1*}

¹ College of Horticulture and Landscape Architecture, Northeast Agricultural University, Harbin 150030, China

² Heilongjiang Forest Botanical Garden, Harbin 150046, China

³ Large-Scale Instrument and Equipment Sharing Service Platform, Northeast Agricultural University, Harbin 150030, China

* Corresponding authors, E-mail: haizhenzhang@neau.edu.cn; aiminzhou@neau.edu.cn

Abstract

Poplar is a tree species used in urban greening, and colored leaf poplar is in great demand because of its higher ornamental value and function. In this study, we generated a golden leaf 84 K poplar (*Populus alba* × *Populus glandulosa*) by overexpressing the *AtSep2* (stress-enhanced protein) gene from *Arabidopsis*. *AtSep2* belongs to the light-harvesting chlorophyll (Chl)-binding complex (LHC)-like family and contains a conserved Chl-binding motif. Subcellular localization analysis showed that *AtSep2* was mainly localized in the chloroplasts of poplar leaves. *AtSep2* overexpression caused the golden leaf phenotype in poplar by reducing total Chl content. Before and after high light stress, the relative reduction in total Chl content and maximum quantum yield of PSII photochemistry (Fv/Fm) in transgenic poplar leaves was lower than that in wild-type (WT) leaves. Transgenic poplar leaves accumulated less reactive oxygen species (ROS) and had a higher net photosynthetic rate (Pn) than WT leaves under high light stress. Overall, *AtSep2* overexpression caused the golden leaf phenotype and enhanced the tolerance of poplar leaves to high light stress. The present study suggests that *AtSep2* is ideal for the genetic improvement of the golden leaf phenotype in poplar and other ornamental plants.

Citation: Chao D, Zhai X, Ren L, Fu S, Zhang J, et al. 2024. Chlorophyll-binding protein *AtSep2* overexpression causes leaf yellowing and enhances tolerance to high light stress in poplar. *Ornamental Plant Research* 4: e018 <https://doi.org/10.48130/opr-0024-0016>

Introduction

Plants with colored leaves are popular in landscaping because of their high ornamental value. Leaf color changes are complex and determined mainly by various pigments in the leaves, such as chlorophyll (Chl), carotenoids (Car), and anthocyanins^[1]. Red or purple leaves are associated with high anthocyanin levels in several woody plants, including *Acer rubrum*^[2], *Populus deltoides*^[3], and *Lagerstroemia indica*^[4]; and similarly golden leaves are associated with low Chl levels in plants, such as *Ulmus pumila*^[5,6], *Ginkgo biloba*^[1], and *Lagerstroemia indica*^[7]. Unlike anthocyanins, Chl affects the leaf color of plants and is a key component of photosynthesis^[8], so its dynamic regulation is more complicated. The Chl metabolic pathway in higher plants consists of the following three steps: Chl biosynthesis, the Chl cycle (interconversion of Chl a and Chl b), and Chl degradation^[9,10]. Inhibition of Chl biosynthesis or acceleration of Chl degradation results in a leaf yellowing phenotype, known as golden leaf. For example, the inhibition of Mg-chelatase subunits I (ChlI) and D (ChlD), which are involved in Chl biosynthesis, causes leaf yellowing in rice and peas (*Pisum sativum*)^[11,12]. The overexpression of staygreen (*SGR*) and pheophytin pheophorbide hydrolase (*PPH*), which regulate Chl degradation, causes a leaf-yellowing phenotype in rice and *Arabidopsis*^[13,14].

In addition to the genes involved in Chl biosynthesis and degradation, the altered expression of light-harvesting Chl-binding complexes (LHC) and LHC-like genes also affect Chl

levels. The conserved Chl-binding motif in the transmembrane helix is a key feature of both LHC and LHC-like proteins^[15]. The Chl bound by the peripheral LHCI and LHCII of photosystems (PS) I and PSII absorbs light energy and transfers this excitation energy to the PS reaction centers. However, LHC-like proteins that bind with free Chl mainly protect PS by dissipating excess energy and participating in the non-photochemical quenching processes^[15]. The LHC-like family comprises three-helix early light-induced proteins (Elips)^[16], two-helix stress-enhanced proteins (Seps), light harvesting-like 3 (Lil3s, also called Sep3s)^[17], and one-helix proteins (Ohps)^[18]. *AtElip2* overexpression causes leaf yellowing in *Arabidopsis* by reducing Chl content^[19]. The *AtSep2* protein contains 2 transmembrane structures, and its conserved Chl-binding site is located in the first transmembrane structure^[20]. Our recent study showed that *AtSep2* overexpression caused leaf yellowing in *Arabidopsis* by reducing Chl content. Moreover, transgenic *Arabidopsis* leaves overexpressing *AtSep2* accumulated less reactive oxygen species (ROS) compared to wild-type (WT) leaves under high-light stress^[20]. These results indicate that *AtSep2* overexpression not only causes leaf yellowing but also reduces oxidative damage in transgenic plants. Therefore, LHC-like genes, especially *AtSep2*, may be used to genetically improve plants to have a golden leaf phenotype and tolerance to high light stress.

Poplars are an important tree species for urban greening, afforestation, and ecological and environmental protection because of their rapid growth, good morphology, and easy

vegetative reproduction. Poplars with the golden leaf poplar (JHY), a bud mutation from *Populus* sp. Linn. '2025'^[21], are economically more valuable because of their higher ornamental value. However, studies on the use of transgenic techniques to generate golden leaf poplar are scarce. In this study, the *AtSep2* gene was overexpressed in the 84 K poplar (*Populus alba* × *Populus glandulosa*), which was then subjected to leaf color and physiological change analysis. Our results showed that *AtSep2* overexpression caused leaf yellowing in 84 K poplar, namely the golden leaf phenotype, indicating that it can be used as a genetic resource for the improvement of golden leaves in ornamental plants.

Methods

Plant materials and growth conditions

Seedlings and plants of the 84 K (*Populus alba* × *Populus glandulosa*) poplar were grown in a greenhouse (24 ± 2 °C) under a 12 h light/12 h dark photoperiod at normal ($150 \mu\text{mol}/\text{m}^2/\text{s}$) or high ($500 \mu\text{mol}/\text{m}^2/\text{s}$) light intensity.

Genetic transformation in the poplar

The open reading frame (ORF) (without the stop codon) of *AtSep2* (At2g21970) (Supplemental Fig. S1) was amplified using the primers *AtSep2*-F (5'-GGTACCATGGCTATGGCTACGCGAG-3') and *AtSep2*-R (5'-GTCGACAAGATCATCAGACCAATCACT-3') and cloned into a pCambia1300-GFP (green fluorescent protein) vector using the *Kpn* I and *Sal* I sites. The reaction components per 20 μL were as follows: 7 μL H_2O , 10 μL Premix Taq™ (RR003A, Takara, Japan), 1 μL 10 μM of each primer and 1 μL cDNA. The thermal cycling program was as follows: initial denaturation at 95 °C for 60 s, and 40 cycles at 95 °C for 10 s, 65 °C for 30 s, and 72 °C for 45 s. The construct was transformed into an 84 K poplar using the *Agrobacterium*-mediated leaf disc method, as described previously^[22–24]. The concentration of *Agrobacterium* carrying the *AtSep2* plasmid was controlled at $\text{OD}_{600} \approx 0.8$ and co-cultured for 25 min. The leaves of 84 K poplar were incubated on 2.41 g/L woody plant media (pH 5.8) supplemented with 0.001 mg/L thidiazuron, 0.06 mg/L 6-benzylaminopurine, 0.01 mg/L indolebutyric acid (IBA), 200 mg/L cefotaxime (Cef), 3 mg/L hygromycin (Hyg), 25 g/L sucrose, and 0.8% (w/v) agar for shoot induction and selection. The regenerated shoots were individually detached from the callus and inserted into a rooting media (Murashige and Skoog [MS] medium supplemented with 0.05 mg/L IBA, 0.02 mg/L 1-naphthaleneacetic acid, 200 mg/L Cef, 3 mg/L Hyg, and 0.9% [w/v] agar). Transgenic poplar plants were identified using reverse transcription PCR. *PtActin* (*PtActin*-F: 5'-GAAGTCCTCTTCCAGCCTTCTC-3'; *PtActin*-R: 5'-CTTGATCTTCATGCTGCTTGGG-3') was used as the internal reference gene.

Chlorophyll (Chl) content and Chl fluorescence parameters

Pigments were extracted from the leaf tissues using 80% ice-cold acetone. Chl a, Chl b, and carotenoids were quantified using a UV/Vis spectrophotometer. The concentration of Chl a, Chl b, and Car were measured separately for light absorption values at 470, 645, and 663 nm, respectively. Chl fluorescence parameters, including minimum Chl fluorescence (F_0), maximum Chl fluorescence (F_m), and maximum quantum yield of PSII photochemistry (F_v/F_m), were measured using kinetic chlorophyll fluorescence imaging systems (FluorCam, PSI,

Czech Republic) at 76.2 μE actinic light and 906 μE saturating pulse.

Subcellular localization

Leaves of transgenic poplars stably expressing *AtSep2*-GFP were cut into small pieces and adhered to glass microscope slides. The GFP signals in the leaf epidermis of poplars were visualized using a confocal laser scanning microscope (Olympus FV3000, Tokyo, Japan). GFP and Chl autofluorescence signals were captured at 488 and 561 nm for excitation and 500–530 and 680–720 nm for emission, respectively.

Transmission electron microscopy

Leaves (0.5 cm × 0.5 cm) of poplars were incubated in the fixation solution (2.5% [v/v] glutaraldehyde in 0.1 M phosphate buffer, pH 7.4) under a vacuum for 12 h. The leaves of the fixed plants were dehydrated in ethanol and embedded in LR White resin (Sigma-Aldrich) for 12 h of polymerization at 50 °C. Ultrathin (60–80 nm) sections of poplar leaves were prepared using a Leica EM UC6 ultramicrotome (Leica, Vienna, Austria). The sections were post-stained with uranyl acetate and lead citrate and observed under an H-7500 transmission electron microscope (Hitachi, Tokyo, Japan) operated at 80 kV.

High light stress test

One-month-old soil-grown transgenic and WT poplar seedlings were treated under continuous high-light conditions ($500 \mu\text{mol}/\text{m}^2/\text{s}$). Five plants were treated for each line. After treatment, leaves at the same position were used to detect Chl fluorescence parameters and ROS content.

Detection of ROS and net photosynthetic rate (P_n)

For superoxide anion ($\text{O}_2^{\cdot-}$) detection, detached leaves were immersed in a 6 mM nitroblue tetrazolium (NBT) solution containing 50 mM sodium phosphate (pH 7.5) for 12 h in the dark. To detect hydrogen peroxide (H_2O_2), detached leaves were immersed in 5 mM 3,3'-diaminobenzidine (DAB) solution containing 10 mM MES (pH 3.8) for 12 h in the dark. H_2O_2 content was measured using a Hydrogen Peroxide Assay Kit (BC3590, Solarbio, China), according to the manufacturer's instructions. The P_n was measured using a Li-6400XT portable photosynthetic system (LI-COR, Lincoln, NE, USA).

Statistical analyses

All experiments were conducted at least in three independent biological and three technical replicates. The data were analyzed using one-way analysis of variance by SPSS, and statistically significant differences were calculated based on Student's t-test, with $p < 0.01$ (**) as thresholds for significance.

Results

AtSep2 overexpression causes Chl reduction and yellowing in poplar leaves

AtSep2 contains two putative transmembrane structures, with a conserved Chl-binding motif (ExxGR) located in the first transmembrane structure (Fig. 1a, b). Glutamic acid (E^{105}) in the Chl-binding motif is essential for Chl binding (Fig. 1c)^[20]. To understand whether *AtSep2* functions in poplar, a genetic transformation was performed using the leaf disc method (Supplemental Fig. S2a). Two transgenic poplar lines were generated and identified using reverse transcription PCR (Supplemental Fig. S2b). Based on phenotypic observations,




Fig. 1 Sequence analysis of AtSep2. (a) Amino acid sequence of AtSep2. (b) Predictive model of the AtSep2 protein structure. (c) Arrowheads indicate the conserved Chl-binding site E¹⁰⁵ amino in the first transmembrane structure (TM).

transgenic poplar seedlings overexpressing *AtSep2* exhibited marked leaf yellowing compared with WT leaves (Fig. 2a). This leaf yellowing phenotype continued in the annual plants and offspring of the cuttings (Supplemental Fig. S3a–f). Transgenic poplar grew slightly slower than WT, but there was no significant difference between them (Fig. 2a, Supplemental Fig. S3a–d). Pigment analysis revealed that the Chl a, Chl b, and Car levels in the leaves of transgenic poplar seedlings were significantly lower than that in the WT leaves (Fig. 2b–d), whereas the ratio of Chl a/b was significantly higher than that in the WT leaves (Fig. 2e). These results suggest that *AtSep2* overexpression may cause leaf yellowing in poplar by reducing the Chl content.

AtSep2 localize in chloroplasts of poplar

Accurate localization of nuclear-encoded chloroplast proteins is a prerequisite for its function; therefore, the subcellular localization of *AtSep2* in tobacco and poplar leaves was observed using GFP as a fusion marker. In tobacco leaves transiently expressing GFP as control, the GFP signals were mainly localized in the nucleus and cytoplasm (Supplemental Fig. S4a). However, in tobacco leaves transiently expressing *AtSep2*-GFP, the GFP signals mainly overlapped with the Chl autofluorescence signals (red) (Supplemental Fig. S4a), indicating that *AtSep2*-GFP was mainly localized in chloroplasts. Moreover, regions of tobacco leaves transiently expressing *AtSep2*-GFP showed yellowing, while regions transiently expressing GFP did not (Supplemental Fig. S4b). In transgenic poplar leaves stably expressing *AtSep2*-GFP, GFP signals were also mainly colocalized with Chl autofluorescence signals. However, no GFP signal was detected in the WT leaves (Fig. 3a). Chloroplast morphology observations showed that the transgenic poplar had

smaller starch grains in the chloroplasts and thinner thylakoids than WT poplar (Fig. 3b). These results indicated that *AtSep2* is mainly localized in the chloroplasts and its overexpression affected chloroplast morphology.

AtSep2 overexpression enhances the tolerance of high light stress in poplar leaves

AtSep2 is involved in light-stress response; therefore, the phenotype and physiology of transgenic poplars overexpressing *AtSep2* and WT were compared under high light stress. After 14 days of high light stress, WT leaves were more purple than transgenic leaves (Fig. 4a, b), which is typical under light-induced stress. Before and after the stress, the Chl a, Chl b, and Car contents in the transgenic poplar leaves were significantly lower than those in the WT leaves, but their relative reduction in the WT leaves (approximately 80%) was greater than that in the transgenic leaves (approximately 60%) (Fig. 4c–e). The detection of Chl fluorescence parameters revealed that transgenic poplar leaves had lower F_0 and F_m and higher F_v/F_m values than WT leaves before and after stress (Fig. 5a–d). Lower F_0 and F_m values in transgenic poplar leaves were associated with their lower Chl content, while higher F_v/F_m values indicate that transgenic poplar leaves have higher intrinsic PSII efficiency compared to WT leaves. Moreover, higher F_v/F_m also indicated that high light stress had less effect on transgenic poplar leaves PSII than WT leaves. NBT staining showed that WT leaves had more staining than transgenic poplar leaves after high light stress (Fig. 6a), whereas DAB staining showed no significant difference between the WT and transgenic poplar leaves after high light stress (Fig. 6b). However, transgenic poplar leaves had significantly lower H_2O_2 accumulation than WT leaves after stress (Fig. 6c). Transgenic poplar leaves had




Fig. 2 Phenotype and pigment content of leaves of the wild-type (WT) and transgenic poplar seedlings overexpressing (OE) *AtSep2*. (a) Comparison of leaf (adaxial and abaxial) color of the WT and transgenic poplar seedlings. The content of (b) chlorophyll (Chl) a, (c) Chl b and (d) carotenoid (car) and the ratio of (e) Chl a and b in the leaves of WT and transgenic poplar seedlings. Asterisks indicate significant differences between WT and transgenic plants (** $p < 0.01$; Student's *t*-test). Error bars represent SE ($n = 3$). Scale bars = 1 cm.




Fig. 3 Subcellular localization of *AtSep2* and its effect on chloroplast ultrastructure in poplar. (a) Confocal images of leaves of wild-type (WT) and transgenic poplar stably expressing *AtSep2*-GFP. GFP fluorescence is green and Chl autofluorescence is red. Yellow signal is obtained by merging the GFP and Chl autofluorescence images. (b) Transmission electron microscopy (TEM) images of chloroplasts of wild-type (WT) and transgenic poplar overexpressing *AtSep2*. Red arrows indicate granum-thylakoid membranes, and asterisks indicate starch granules. Scale bars = (a) 20 μ m, (b) 500 nm.

significantly lower *Pn* than WT leaves under normal light conditions. However, *Pn* in transgenic poplar leaves was significantly

higher than that in WT leaves after high light stress (Fig. 6d). Overall, although the transgenic poplar leaves had less Chl under normal conditions, their relative reduction after light stress was also small. Moreover, after light stress, transgenic poplar leaves showed higher *Fv/Fm* and *Pn* than WT leaves. These results indicate that transgenic poplar leaves have better tolerance to high light conditions than WT poplar leaves, as reflected by the higher *Fv/Fm* and *Pn* and lower accumulation of O_2^- and H_2O_2 after high light stress.

Discussion

Plants with differently colored leaves usually arise from natural mutations in their buds, which are stochastic and uncertain. Genetic engineering is an effective way to change the leaf color of plants. Poplar leaf color can be improved using transgenic methods. For example, overexpression of the R2R3-MYB transcription factors *PtrMYB119* and *PdMYB118* caused red leaf phenotypes in *P. trichocarpa* and *P. deltoids*, respectively, by promoting anthocyanin accumulation^[3,25,26]. Suppression of *GLK* (*golden2-like*) gene expression caused the golden leaf phenotype in triploid poplar (*P. alba* \times *P. berlinensis*)^[27]. In the present study, *AtSep2* overexpression caused the golden leaf phenotype in 84 K poplar by reducing Chl levels (Fig. 2). A previous study by the current authors confirmed that *AtSep2* can bind free Chl under light stress^[20]. High light stress causes saturation of the energy absorbed by the PSs. Excess energy triggers the transition of free Chl into Chl triplets, which then react with O_2 to form harmful ROS in the chloroplast^[15]. Thus, less accumulation of free Chl under light stress might be closely associated with less ROS production. In the present study, the transgenic poplar leaves had lower Chl content and less O_2^- and H_2O_2 accumulation than the WT leaves after high light stress (Figs 4, 6). Moreover, transgenic poplar leaves showed a stronger high light tolerance than WT leaves (Figs 4, 5). These results suggest that *AtSep2* reduces ROS production by




Fig. 4 Phenotype and pigment content of the leaves of wild-type (WT) and transgenic poplar overexpressing (OE) *AtSep2* under high light stress. (a) Plant and leaf phenotypes of WT and transgenic poplar under high light stress for 14 d. The content of (b) chlorophyll (Chl) a, (c) Chl b and (d) carotenoid (car) and the ratio of (e) Chl a and b in the leaves of WT and transgenic poplar. Asterisks indicate significant differences between WT and transgenic plants (** $p < 0.01$; Student's t -test). Error bars represent SE ($n = 3$). Scale bars = 1 cm.




Fig. 5 Chlorophyll (Chl) fluorescence parameters in the leaves of wild type (WT) and transgenic poplar overexpressing (OE) *AtSep2* under high light stress. (a) Minimum Chl fluorescence (F_0), maximum Chl fluorescence (F_m), and maximal quantum yield of PSII photochemistry (F_v/F_m) images of the leaves of WT and transgenic poplar under normal light or high light stress for 14 d and (b)–(d) their quantitative analysis. Pseudo-colored bar depicted at the bottom of the panel ranges from 0 (black) to 300 (F_0), 1,000 (F_m), or 1.0 (F_v/F_m) (red). Asterisks indicate significant differences between WT and transgenic plants (** $p < 0.01$; Student's t -test). Error bars represent SE ($n = 3$).

binding more free Chl, thereby enhancing the high light tolerance of transgenic poplar leaves (Fig. 7). Studies have revealed that LHC-like proteins are mainly responsible for photoprotection by reducing light-induced oxidative damage in chloroplasts^[15]. In agreement, a previous study by the current authors also confirmed the function of *AtSep2* in Arabidopsis^[20], suggesting that it plays the same role in both Arabidopsis and poplar. Subcellular localization showed that *AtSep2* can accurately localized to the chloroplasts of tobacco and poplar leaves by transient and stable expression

experiments (Fig. 3a, Supplemental Fig. S4a). Chloroplast proteins are involved in post-transcriptional regulation, and their accurate localization is crucial for their function. The Chl metabolism process is mainly performed in chloroplasts, and the accurate localization of *AtSep2* in chloroplasts also indicates that its function is conserved in poplar leaves.

Due to its higher ornamental value, *AtSep2*-transgenic golden leaf poplar should be more popular in urban greening and beautification. In addition, high-light stress is one of the environmental factors limiting the application of landscaping




Fig. 6 Reactive oxygen species (ROS) accumulation and net photosynthetic rate (P_n) in the leaves of wild type (WT) and transgenic poplar overexpressing (OE) *AtSep2* under high light stress. The O_2^- and H_2O_2 levels in the leaves of WT and transgenic poplar under normal light or high light stress for 14 d, as indicated by (a) nitroblue tetrazolium (NBT) and (b) 3,3'-diaminobenzidine (DAB) staining. (c) H_2O_2 content and (d) P_n in the leaves of WT and transgenic poplar under normal light or high light stress for 14 d. Asterisks indicate significant differences between WT and transgenic plants (** $p < 0.01$; Student's t -test). Error bars represent SE ($n = 3$).




Fig. 7 Putative model of *AtSep2* function in leaf yellowing and tolerance of high light stress in transgenic poplar. High light stress induces free chlorophyll (Chl) to form the Chl triplets, which react with O_2 to form harmful reactive oxygen species (ROS), such as H_2O_2 and O_2^- ; however, *AtSep2* may prevent the further production of ROS by binding with free Chl.

tree species in many areas at present, therefore, *AtSep2*-transgenic poplar with stronger high light tolerance will have greater application prospects. Although *AtSep2* is derived from the herb *Arabidopsis*, the golden leaf phenotype is stable in

transgenic poplar overexpressing *AtSep2* (Figs 2, 4, Supplemental Fig. S3). Moreover, the growth of transgenic golden leaf poplar was slightly inhibited but did not produce a significant difference from WT (Figs 2, 4, Supplemental Fig. S3). Therefore, *AtSep2* is the candidate gene in tree breeding for the generation of species with different leaf colors.

Conclusions

In summary, *AtSep2* has a conserved Chl-binding motif, and its overexpression causes the golden leaf phenotype in poplar by reducing Chl content. *AtSep2* was mainly localized in the chloroplasts in the leaves, and its overexpression altered the chloroplast micromorphology. Under high light stress, the transgenic poplar leaves exhibited higher light tolerance than the WT leaves, as indicated by a lower decrease in F_v/F_m , ROS accumulation, and higher P_n . Overall, *AtSep2* may be used for the genetic improvement of golden leaf poplar through genetic engineering.

Author contributions

The authors confirm contribution to the paper as follows: literature collation: Chao D, Zhou A; experimental design and implementation: Zhang H, Zhou A; data analysis and visualization: Chao D, Zhai X, Ren L; writing and polishing: Chao D, Fu S, Zhang J; resources and project administration: Zhai X, Feng S;

review and editing: Zhang H, Zhou A. All authors reviewed the results and approved the final version of the manuscript.

Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (32172619).

Conflict of interest

The authors declare that they have no conflict of interest.

Supplementary information accompanies this paper at (<https://www.maxapress.com/article/doi/10.48130/opr-0024-0016>)

Dates

Received 20 March 2024; Revised 11 May 2024; Accepted 3 June 2024; Published online 2 July 2024

References

- Li W, Yang S, Lu Z, He Z, Ye Y, et al. 2018. Cytological physiological and transcriptomic analyses of golden leaf coloration in *Ginkgo biloba* L. *Horticulture Research* 5:12
- Chen Z, Lu X, Xuan Y, Tang F, Wang J, et al. 2019. Transcriptome analysis based on a combination of sequencing platforms provides insights into leaf pigmentation in *Acer rubrum*. *BMC Plant Biology* 19:240
- Wang H, Wang X, Song W, Bao Y, Jin Y, et al. 2019. PdMYB118 isolated from a red leaf mutant of *Populus deltoides* is a new transcription factor regulating anthocyanin biosynthesis in poplar. *Plant Cell Reports* 38:927–36
- Feng L, Shen P, Chi X, Zhou Y, Liu J, et al. 2023. The anthocyanin formation of purple leaf is associated with the activation of *LfiHY5* and *LfiMYB75* in crape myrtle. *Horticultural Plant Journal* 10:1230–46
- Zhang S, Zuo L, Zhang J, Chen P, Wang J, et al. 2017. Transcriptome analysis of *Ulmus pumila* 'Jinye' responses to different shading involved in chlorophyll metabolism. *Tree Genetics & Genomes* 13:64
- Zuo L, Zhang S, Liu Y, Huang Y, Yang M, et al. 2019. The reason for growth inhibition of *Ulmus pumila* 'Jinye': lower resistance and abnormal development of chloroplasts slow down the accumulation of energy. *International Journal of Molecular Sciences* 20:4227
- Li S, Wang S, Wang P, Gao L, Yang R, et al. 2020. Label-free comparative proteomic and physiological analysis provides insight into leaf color variation of the golden-yellow leaf mutant of *Lagerstroemia indica*. *Journal of Proteomics* 228:103942
- Wang P, Richter AS, Kleeberg JRW, Geimer S, Grimm B. 2020. Post-translational coordination of chlorophyll biosynthesis and breakdown by BCMs maintains chlorophyll homeostasis during leaf development. *Nature Communications* 11:1254
- Mochizuki N, Tanaka R, Grimm B, Masuda T, Moulin M, et al. 2010. The cell biology of tetrapyrroles: a life and death struggle. *Trends in Plant Science* 15:488–98
- Woo HR, Kim HJ, Lim PO, Nam HG. 2019. Leaf senescence: systems and dynamics aspects. *Annual Review of Plant Biology* 70:347–76
- Luo T, Luo S, Araújo WL, Schlicke H, Rothbart M, et al. 2013. Virus-induced gene silencing of pea *CHL1* and *CHLD* affects tetrapyrrole biosynthesis, chloroplast development and the primary metabolic network. *Plant Physiology and Biochemistry* 65:17–26
- Zhang H, Li J, Yoo JH, Yoo SC, Cho SH, et al. 2006. Rice *Chlorina-1* and *Chlorina-9* encode ChlD and ChlI subunits of Mg-chelatase, a key enzyme for chlorophyll synthesis and chloroplast development. *Plant Molecular Biology* 62:325–37
- Park SY, Yu JW, Park JS, Li J, Yoo SC, et al. 2007. The senescence-induced staygreen protein regulates chlorophyll degradation. *The Plant Cell* 19:1649–64
- Schelbert S, Aubry S, Burla B, Agne B, Kessler F, et al. 2009. Pheophytin pheophorbide hydrolase (pheophytinase) is involved in chlorophyll breakdown during leaf senescence in *Arabidopsis*. *The Plant Cell* 21:767–85
- Levin G, Schuster G. 2023. LHC-like proteins: the guardians of photosynthesis. *International Journal of Molecular Sciences* 24:2503
- Montané MH, Kloppstech K. 2000. The family of light-harvesting-related proteins (LHCs, ELIPs, HLIPs): was the harvesting of light their primary function? *Gene* 258:1–8
- Heddad M, Adamska I. 2000. Light stress-regulated two-helix proteins in *Arabidopsis thaliana* related to the chlorophyll *a/b*-binding gene family. *Proceedings of the National Academy of Sciences of the United States of America* 97:3741–46
- Andersson U, Heddad M, Adamska I. 2003. Light stress-induced one-helix protein of the chlorophyll *a/b*-binding family associated with photosystem I. *Plant Physiology* 132:811–20
- Tzvetkova-Chevolleau T, Franck F, Alawady AE, Dall'Osto L, Carrière F, et al. 2007. The light stress-induced protein ELIP2 is a regulator of chlorophyll synthesis in *Arabidopsis thaliana*. *The Plant Journal* 50:795–809
- Ren L, Ma H, Chao D, Zhang H, Qiao K, et al. 2023. Sep2, a light-harvesting complex-like protein, is involved in light stress response by binding to free chlorophylls. *Environmental and Experimental Botany* 213:105429
- Tian Y, Rao S, Li Q, Xu M, Wang A, et al. 2021. The coloring mechanism of a novel golden variety in *Populus deltoides* based on the RGB color mode. *Forestry Research* 1:5
- He F, Wang HL, Li HG, Su Y, Li S, et al. 2018. PeCHYR1, a ubiquitin E3 ligase from *Populus euphratica*, enhances drought tolerance via ABA-induced stomatal closure by ROS production in *Populus*. *Plant Biotechnology Journal* 16:1514–28
- Hu X, Wang S, Zhang H, Zhang H, Feng S, et al. 2022. Plant cadmium resistance 6 from *Salix linearistipularis* (SIPCR6) affects cadmium and copper uptake in roots of transgenic *Populus*. *Ecotoxicology and Environmental Safety* 245:114116
- Yao W, Wang S, Zhou B, Jiang T. 2016. Transgenic poplar overexpressing the endogenous transcription factor *ERF76* gene improves salinity tolerance. *Tree Physiology* 36:896–908
- Cho JS, Nguyen VP, Jeon HW, Kim MH, Eom SH, et al. 2016. Overexpression of *PtMYB119*, a R2R3-MYB transcription factor from *Populus trichocarpa*, promotes anthocyanin production in hybrid poplar. *Tree Physiology* 36:1162–76
- Wang H, Wang X, Yu C, Wang C, Jin Y, et al. 2020. MYB transcription factor *PdMYB118* directly interacts with bHLH transcription factor *PdTT8* to regulate wound-induced anthocyanin biosynthesis in poplar. *BMC Plant Biology* 20:173
- Li Y, Gu C, Gang H, Zheng Y, Liu G, et al. 2021. Generation of a golden leaf triploid poplar by repressing the expression of *GLK* genes. *Forestry Research* 1:3



Copyright: © 2024 by the author(s). Published by Maximum Academic Press, Fayetteville, GA. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.