

Molecular taxonomy reveals new records of *Chromolaenicola* (Didymosphaeriaceae, Pleosporales) and potential antibacterial properties

Zin Hnin Htet^{1,2}, Ausana Mapook^{2*} and K. W. Thilini Chethana^{1,2*}

¹ School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

² Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand

* Corresponding authors, E-mail: phung.ausana@gmail.com; kandawatte.thi@mfu.ac.th

Abstract

In the present study, *Chromolaenicola* specimens were collected from the dead stems of *Bidens pilosa* in northern Thailand and compared with other *Chromolaenicola* species. Maximum-likelihood and Bayesian analyses were conducted on a combined LSU, SSU, ITS, *tef-1*, and *rpb2* data set. Morphological and phylogenetic analyses revealed three species, *Chromolaenicola chiangraiensis* (new host record), *C. siamensis* (new host record), and *C. thailandensis* (new host and asexual morph record), respectively. A preliminary screening demonstrated the ability of *Chromolaenicola chiangraiensis*, *C. thailandensis*, and *C. siamensis* to partially inhibit the growth of gram-positive bacteria (*Bacillus subtilis*). Here, the detailed morphology, illustrations, and potential antibacterial properties of *Chromolaenicola* species are described. The present research will add to the body of knowledge on *Chromolaenicola* by revealing its possible antibacterial properties.

Citation: Htet ZH, Mapook A, Chethana KWT. 2024. Molecular taxonomy reveals new records of *Chromolaenicola* (Didymosphaeriaceae, Pleosporales) and potential antibacterial properties. *Studies in Fungi* 9: e006 <https://doi.org/10.48130/sif-0024-0006>

Introduction

As a major driver of biodiversity loss, invasive plant species threaten the natural environment and human health globally^[1–7]. Invasive plant species not only disrupt natural community assembly but also act as one of the most significant impediments to restoring native ecosystems^[7,8]. *Bidens pilosa* is a widespread weed in tropical, subtropical and warm temperate regions^[9]. This weed is a diverse annual herb native to tropical and Central America^[9] and is considered an invasive weed in Thailand^[10]. Even though several studies have been conducted on fungi associated with *Bidens pilosa*^[11–13], the comprehensive fungal diversity of *Bidens pilosa* is still yet to be understood.

Fungi associated with *Bidens pilosa* have been observed in some studies, and the reported species belong to the families Albuginaceae, Apiosporaceae, Botryosphaeriaceae, Ceratobasidiaceae, Cladosporiaceae, Diaporthaceae, Erysiphaceae, Glomerellaceae, Mycosphaerellaceae, Nectriaceae, Periconiaceae, Peronosporaceae, Phyllostictaceae, Sclerotiniaceae, Stachybotryaceae, Tetraplosporaaceae, and Torulaceae^[11–14]. However, species of Didymosphaeriaceae have not been reported from *Bidens pilosa* yet. Didymosphaeriaceae, as introduced by Munk^[15], comprises 33 genera^[16]. Members of this family can be found as endophytes, pathogens and saprobes in soil as well as various host plants from both aquatic and terrestrial habitats^[17–24]. Among them, *Chromolaenicola*, which was introduced by Mapook et al.^[17], accommodated seven accepted species^[16,25,26], including four asexual morph (*Chromolaenicola chiangraiensis*, *C. clematidis*, *C. lampangensis*, and *C. siamensis*)

and three sexual morph taxa (*C. nanensis*, *C. sapinda* and *C. thailandensis*)^[17,27]. *Chromolaenicola* species occur as saprobes in terrestrial habitats in China and Thailand^[17,27–29]. Members of *Chromolaenicola* were found from different plant families, viz., Asteraceae, Bromeliaceae, Fabaceae, Ranunculaceae, and Spinaceae^[17,27–30]. Some *Chromolaenicola* species, such as *C. lampangensis* and *C. nanensis*, showed potential antimicrobial properties^[17]. The sexual morph of *Chromolaenicola* is characterized by its immersed to semi-immersed, solitary or scattered, coriaceous, globose to subglobose ascomata with an ostiole, 3–4 layers of *textura angularis* cells, cylindrical to filiform, septate, branching pseudoparaphyses, 6–8-spored, bitunicate, cylindrical, pedicellate asci with an ocular chamber, and hyaline to brown, uniseriate, ellipsoid to broadly fusiform, muriform spores without gelatinous sheath. The asexual morph is characterized by its immersed to semi-immersed, globose to obpyriform, pycnidia conidiomata with an ostiole, 2–4 layers of *textura angularis* cells, hyaline and unbranched, smooth, elongated, broadly filiform to ampulliform conidiogenous cells, and oblong or oval to ellipsoid, globose to subglobose, hyaline to pale brown, aseptate to 1-septate conidia. The linkage of sexual and asexual morph from *Chromolaenicola* has not been reported yet.

In the current study, morphological descriptions and photo plates of *Chromolaenicola* species collected in northern Thailand are presented. Both multi-gene analyses and morphological comparisons were used to confirm the species identification. The preliminary antibacterial screening was also conducted and presented for all *Chromolaenicola* species.

Materials and methods

Sample collection and morphological study and isolation

Dead stems of *Bidens Pilosa* were collected in March from the roadsides in Doi Pui District, Chiang Rai Province, Thailand. All specimens were kept in plastic bags labeled with collection details and taken to the laboratory. Single spore isolation was carried out on malt extract agar (MEA), following the methods by Senanayake et al.^[31] and kept for 24 h at room temperature. The spore germination was observed within 24 h using a Motic SMZ 168 Series microscope. Germinated spores were transferred to new MEA plates. All the micro-morphological characters were observed using a Nikon ECLIPSE 80i compound microscope (Nikon, Japan) fitted to a Canon 550D digital camera (Canon, Japan). Tarosoft Image Framework (v 0. 9.7) was used to measure photomicrograph structures. Adobe Photoshop CS6 Extended (v 10.0.) was used to edit and prepare photo plates (Adobe Systems, USA). Forty-day-old cultures were used for molecular studies. Specimens were deposited at the herbaria of Mae Fah Luang University (Herb. MFLU) while living cultures were maintained at Mae Fah Luang University culture collection (MFLUCC). Faces of fungi (FoF) numbers and Index Fungorum (IF) numbers were obtained as instructed by Jayasiri et al.^[32] and Index fungorum^[25]. Moreover, the species descriptions were submitted to GMS Microfungi^[33].

DNA extraction, PCR amplification and sequencing

Genomic DNA was extracted from forty-day-old mycelium using E.Z.N.A.® Tissue DNA Kit (Omega Biotek Inc.), following the manufacturer's instructions. DNA amplifications were performed by polymerase chain reactions (PCR), following Mapook et al.^[34]. The details of PCR primers and protocols are shown in Table 1. The quality of PCR products was confirmed on 1% gels. The PCR products were sent to a commercial sequencing provider (Solgent Co., Ltd, Thailand). The newly generated nucleotide sequences were deposited in GenBank, and accession numbers were obtained (Table 2).

Sequence alignment and phylogenetic analyses

Newly generated forward and reverse sequences were assembled in the SeqMan^[47]. The assembled sequences were used for BLAST searches at NCBI (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). Based on previously published data^[17,27,30] and BLAST search results, taxa were selected, and phylogenetic analyses were conducted using the combined LSU, SSU, ITS, *tef1-α*, and *rpb2* sequence data. Sequence alignments were made with the MAFFT v. 7 online tool (<http://mafft.cbrc.jp/alignment/server>; 2016). Using MEGA v. 6.0, alignments were improved where necessary, and composite sequence alignments were obtained^[48].

Maximum likelihood (ML) and Bayesian inference (BI) analyses were conducted using the combined dataset of LSU, SSU, ITS, *tef1-α*, and *rpb2*^[49,50]. RA × ML and Bayesian analyses were carried out on the CIPRESS Science Gateway Portal (www.phylo.org) using the methods described by Miller et al.^[51]. Maximum likelihood analysis was performed by RA × ML-HPC v.8^[52] with rapid bootstrap analysis, followed by 1000 bootstrap replicates and the GTRGAMMA substitution model. MrBayes was used to perform BI analysis on XSEDE 3.2.7^[53], with tree samples taken at every 100th generation during the 5M generation run of four concurrent Markov chains. The first 25% of the trees were removed as part of the burn-in phase, and calculations for the Posterior Probability were made for the remaining 75% of the trees (PP)^[49,54]. The phylogenetic tree was displayed using Fig Tree v1.4.0^[55] and was modified in Microsoft Office PowerPoint v. 2013.

Preliminary screening for antibacterial activity

Preliminary screening for antimicrobial activity was carried out following the methods of Mapook et al.^[17]. Antibacterial discs of Ampicillin were used as a positive control for screening^[56]. Preliminary antibacterial activities were tested against *Bacillus subtilis* (TISTR 1248), *Escherichia coli* (TISTR 527), and *Staphylococcus aureus* (TISTR Y4b) using the agar plug diffusion method^[57]. Bacterial test organisms were grown on Nutrient Agar (NA) for 24 h. After 24 h of inoculation, 2–3 loops of the bacterial test organisms were transferred to the nutrient broth. Before adding microbial suspensions to the sterile Mueller-Hinton agar media, cell counts were performed on the suspensions (6.7×10^5 cells/mL), as detailed by Mapook et al.^[17]. Fungal mycelium plugs from our isolates were transferred to a solid medium plate and allowed to grow at room temperature for 24–48 h. Inhibition zones were measured and compared to the positive control.

Results

Phylogenetic analyses

Phylogenetic analysis was conducted from the combined LSU, SSU, ITS, *tef1-α*, and *rpb2* sequence data of twenty-eight strains, of which three were newly sequenced, while the other 23 strains were obtained from BLAST search (NCBI) and recent papers^[17,27,29,30]. *Periconia pseudodigitata* strains KT1395 and KT1195A were selected as the outgroup. The best-scoring RA×ML tree with a final likelihood value of −9277.475085 is presented (Fig. 1). The matrix had 532 distinct alignment patterns, with 15.55% of undetermined characters or gaps. Estimated base frequencies were as follows: A = 0.235845, C = 0.255030, G = 0.269910, T = 0.239215; substitution rates: AC = 1.648158, AG = 2.210120, AT = 1.275667, CG = 1.033223, CT = 7.205216, GT = 1.000000; gamma distribution shape parameter α = 0.110629.

Table 1. PCR conditions used in this study.

Gene	Primers		PCR conditions	Ref.
	Forward	Reverse		
Large subunit (LSU)	LR0R	LR5	95 °C: 3 min, (94 °C: 30 s, 56 °C: 50 s, 72 °C: 1 min) × 40 cycles 72 °C: 7 min	[35]
Small subunit (SSU)	NS1	NS4	95 °C for 3 min, (94 °C: 30 s, 55 °C: 50 s, 72 °C: 1 min) × 40 cycles 72 °C: 7 min	[36]
Internal transcribed spacer (ITS)	ITS5	ITS4	95 °C for 3 min, (94 °C: 30 s, 55 °C: 50 s, 72 °C: 1 min) × 40 cycles 72 °C: 7 min	[36]
Elongation factor-1 alpha (<i>tef1-α</i>)	EF-1 983F	EF1-2218R	95 °C: 3 min, (94 °C: 30 s, 55 °C: 50 s, 72 °C: 1 min) × 40 cycles 72 °C: 7 min	[37]
RNA polymerase II subunit (<i>rpb2</i>)	frPB2-5 F	frPB2-7cR	95 °C: 5 min, (95 °C: 1 min, 52 °C: 2 min, 72 °C: 90 s) × 40 cycles 72 °C: 10 min	[38]

Table 2. List of taxa, specimens and sequences used in phylogenetic analyses.

Species	Strain numbers	GenBank accession numbers					Ref.
		LSU	SSU	ITS	<i>tef1-α</i>	<i>rpb2</i>	
<i>Bimuria omanensis</i>	SQUCC 15280	NG_071257	N/A	NR_173301	MT279046	N/A	[39]
<i>B. novae-zelandiae</i>	CBS 107.79	MH872950	NA	MH861181	NA	N/A	[40]
<i>Chromolaenicola ananasi</i>	MFLU 23-0167	OR438811	OR458332	OR438340	OR500305	N/A	[30]
<i>C. clematidis</i>	MFLUCC 17-2075 ^T	MT310601	MT214554	MT226671	N/A	N/A	[29]
<i>C. chiangraiensis</i>	MFLUCC 17-1493	MN325005	MN325011	MN325017	MN335650	MN335655	[17]
<i>C. chiangraiensis</i>	MFLUCC 24-0058	PP464125	PP464129	PP464138	PP474193	PP474190	This study
<i>Chromolaenicola nanensis</i>	MFLUCC 17-1477	MN325002	MN325008	MN325014	MN335647	MN335653	[17]
<i>C. nanensis</i>	MFLUCC 17-1473	MN325003	MN325009	MN325015	MN335648	MN335653	[17]
<i>C. lampangensis</i>	MFLUCC 17-1462	MN325004	MN325010	MN325016	MN335649	MN335654	[17]
<i>C. siamensis</i>	MFLUCC 17-2527	NG_066311	N/A	NR_163337	N/A	N/A	[28]
<i>C. siamensis</i>	MFLUCC 24-0057	PP464124	PP464128	PP464137	PP474192	PP474189	This study
<i>C. sapindi</i>	KUMCC 21-0564 ^T	OP059009	OP059058	OP058967	OP135943	N/A	[27]
<i>C. sapindi</i>	KUMCC 21-0594	OP059010	OP059059	OP058968	OP135944	N/A	[27]
<i>C. thailandensis</i>	MFLUCC 17-1510	MN325006	MN325012	MN325018	MN335651	N/A	[17]
<i>C. thailandensis</i>	MFLUCC 17-1475	MN325007	MN325013	MN325019	MN335652	MN335656	[17]
<i>C. thailandensis</i>	MFLUCC 24-0056	PP464123	PP464127	PP464136	PP474191	PP474188	This study
<i>Deniquelata barringtoniae</i>	MFLUCC 11-0422	JX254655	JX254656	NR_111779	N/A	N/A	[41]
<i>D. quercina</i>	ABRIICC 10068	MH316157	MH316155	MH316153	N/A	N/A	[42]
<i>Didymocrea leucaenae</i>	MFLUCC 17-0896	NG_066304	MK347826	NR_164298	MK360052	N/A	[28]
<i>D. sadasivanii</i>	CBS 438.65	DQ384103	DQ384066	MH870299	N/A	N/A	[40]
<i>Letendreaa cordylinicola</i>	MFLUCC 11-0148	NG_059530	NG_068362	NR_154118	N/A	N/A	[41]
<i>L. helminthicola</i>	CBS 884.85	AY016362	AY016345	MK404145	MK404174	N/A	[43]
<i>Montagnula acaciae</i>	MFLUCC 18-1636	ON117298	ON117267	ON117280	ON158093	N/A	[44]
<i>M. acaciae</i>	NCYUCC 19-0087	ON117299	ON117268	ON117281	ON158094	N/A	[44]
<i>M. aloes</i>	CPC 19671	JX069847	N/A	JX069863	N/A	N/A	[45]
<i>M. aloes</i>	CBS 132531	NG_042676	N/A	NR_111757	N/A	N/A	[40]
<i>Periconia pseudodigitata</i>	KT1395	AB807564	AB797274	LC014591	AB808540	N/A	[46]
<i>P. pseudodigitata</i>	KT1195A	AB807563	AB797273	LC014590	AB808539	N/A	[46]

^T: Type strains; Abbreviations of culture collections: CBS: Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands, CPC: Working collection of Pedro Crous housed at CBS, KT: K. Tanaka, MFLUCC: Mae Fah Luang University Culture Collection, Chiang Rai, Thailand, NCYUCC: National Chiayi University Culture Collection, Taiwan. SQUCC: Sultan Qaboos University Culture Collection, Sultanate of Oman. ABRIICC: Agricultural Biotechnology Research Institute of Iran Culture Collection, Iran. Sequences generated in the current study are in bold. N/A: Not available.

Tree topologies of ML and BI criteria were comparable to and consistent with earlier investigations^[17,27]. *Chromolaenicola* formed an independent topmost clade in the phylogenetic tree. *Chromolaenicola* species were divided into five clades (Clades A, B, C, D and E) in the phylogenetic tree. Our three strains, represented by three species, were grouped within the genus (Fig. 1). *Chromolaenicola chiangraiensis* (MFLUCC 24-0058) and *C. siamensis* (MFLUCC 24-0057) were placed in Clade A. *Chromolaenicola chiangraiensis* (MFLUCC 24-0058) formed a separate branch with 81% ML and 0.98 BYPP support. Our strain, *Chromolaenicola siamensis* (MFLUCC 24-0057) clustered with *C. siamensis* (MFLUCC 17-2527) with 55% ML and 0.81 BYPP. Three *Chromolaenicola thailandensis* strains (MFLUCC 17-1510, MFLUCC 17-1475, MFLUCC 24-0056) clustered together with 94% ML and 0.99 BYPP support and formed a distinct lineage in Clade C.

Taxonomy

Didymosphaeriaceae Boonmee and K.D. Hyde, Fungal Diversity 80: 462 (2016).

Chromolaenicola chiangraiensis Mapook & K.D. Hyde *Fungal Diversity* 101, 1–175 (2020). (Fig. 2).

Index fungorum number: IF557280, *Faces of fungi number*: FOF 07784

Saprobic on dead stems of *Bidens pilosa*. **Sexual morph**: Undetermined. **Asexual morph**: appearing as *Colonies* on the host substrate, superficial, scattered, gregarious, black. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* 2–4 × 1–2 μm (\bar{x} = 2.5 × 1.6 μm, n = 10), holoblastic, hyaline, smooth, ovoid to filiform. *Conidia* 9–13 × 6–10 μm (\bar{x} = 10.1 × 6.9 μm, n = 30), oval to slightly ellipsoidal, aseptate when immature, 1-septate when mature, thick-walled, reddish brown, verruculose.

Culture characteristics: Conidia germinating on MEA within 24 h, reaching 20 mm after 10 d at room temperature, irregular, undulate, curled margin, yellow to pale brown on the surface and wrinkle and brown in reverse.

Material examined: Thailand, Chiang Rai Province, Doi Pui, on dead stems of *Bidens pilosa*, 14 March 2023, Zin Hnin Htet (BP-DP-10, MFLU 24-0030, **new host record**); living culture MFLUCC 24-0058.

Notes: Morphologically, our species, *Chromolaenicola chiangraiensis* (MFLUCC 24-0058) is similar to *C. chiangraiensis* (MFLUCC 17-1493) in having superficial, scattered, dark brown




Fig. 1 Phylogram generated from maximum likelihood analysis based on the combined dataset of LSU, SSU, ITS, *tef1-α* and *rpb2* sequence data. Bootstrap support values for ML equal to or greater than 75% and BYPP equal to or greater than 0.95 are given at the nodes. Newly generated sequences are in blue and type species are in bold.

to black colonies, holoblastic, hyaline conidiogenous cells, oval to ellipsoidal, aseptate to 1-septate, reddish brown, verruculose conidia but differ in having shorter conidiogenous cells (2–4 vs 3.5–6.5 μm) (Table 3). A comparison of the *tef1-α* gene region of *Chromolaenicola Chiangraiensis* (MFLUCC 24-0058) and *C. Chiangraiensis* (MFLUCC 17-1493) reveals 13 base pair differences (1.48%) across 876 nucleotides. Therefore, our strain (MFLUCC 24-0058) is described as a new host record of *Chromolaenicola Chiangraiensis* from *Bidens pilosa* (Asteraceae), which was previously recorded from *Chromolaena odorata* (Asteraceae) in Chiang Rai Province, Thailand^[17].

Chromolaenicola siamensis (Jayasiri, E.B.G. Jones & K.D. Hyde) Mapook & K.D. Hyde *Fungal Diversity* **101**, 1–175 (2020). (Fig. 3).

Index fungorum number: IF557283, *Faces of fungi* number: FOF 07787

Saprobic on dead stems of *Bidens pilosa*. **Sexual morph:** Undetermined. **Asexual morph:** *Conidiomata* 130–235 × 170–230 μm (\bar{x} = 150.6 × 225.5 μm, *n* = 5), pycnidial, solitary, immersed to semi-immersed, uniloculate, globose, yellowish

brown to brown, without ostiole. *Peridium* 15–20 μm wide, comprising 2–3 layers of brown cells of *textura angularis*. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* 1–2.5 × 2–4 μm (\bar{x} = 2 × 2.5 μm, *n* = 30), phialidic, hyaline. *Conidia* 7–15 × 5–10 μm (\bar{x} = 10.4 × 6.7 μm, *n* = 30), globose to subglobose, 1-septate, thick-walled, reddish brown to dark brown, verruculose.

Culture characteristics: Conidia germinating on MEA after 24 h, 22 mm after 10 d at room temperature, irregular, entire, curled margin, yellow to pale brown on the surface, wrinkled and brown in reverse.

Material examined: Thailand, Chiang Rai Province, Doi Pui, on dead stems of *Bidens pilosa*, 14 March 2023, Zin Hnin Htet (BP-DP-7, MFLU 24-0029, **new host record**); living culture MFLUCC 24-0057.

Notes: In our phylogenetic study, our strain (MFLUCC 24-0057) is sistered to *Chromolaenicola siamensis* (MFLUCC 17-1527) with 51% ML and 0.82 BYPP. When we compared the morphology, our strain (MFLUCC 24-0057) is similar to




Fig. 2 *Chromolaenicola chaingraiensis* (MFLU 24-0030, new host record). (a), (b) Colonies on the substrate. (c)–(f) Conidia and conidiogenous cells. (g)–(j) Conidia. (k) Germinating conidia. (l) Culture on MEA. Scale bars: (a), (b) = 500 μ m, (c) = 30 μ m, (d)–(j) = 10 μ m, (k) = 20 μ m, (l) = 10 mm.

Chromolaenicola siamensis (MFLUCC 17-1527) in having pycnidial, solitary, immersed, globose to obpyriform, unilocular conidiomata, phialidic, hyaline conidiogenous cells, and hyaline to dark brown, globose to subglobose, aseptate to 1-septate conidia with similar size (7–15 \times 5–10 μ m vs 7.2–9.4 \times 5.4–6.5 μ m). Our strain (MFLUCC 24-0057) differs from *C. siamensis* (MFLUCC 17-1527) in having thinner peridium (15–20 μ m vs 15–38 μ m) and shorter conidiogenous cells (1–2.5 \times 2–4 μ m vs 6.5–7.4 \times 3.2–4.7 μ m) (Table 3). However, the comparison of base pair differences revealed no or insignificant results LSU 0% (0/851), ITS 0.2% (1/459), *tef1*- α 0.3% (2/740), *rpb2* 0.1% (1/914), which indicates that they are conspecific. Therefore, we herein report *C. siamensis* as a new host record from *Bidens Pilosa* (Asteraceae), while this species was previously recorded on the decaying pods of *Leucaena* sp. (Fabaceae) in Lampang Province, Thailand[28].

Chromolaenicola thailandensis Mapook & K.D. Hyde *Fungal Diversity* **101**, 1–175 (2020). (Fig. 4).

Index fungorum number: IF557284, *Faces of fungi number*: FOF 07788

Saprobic on the dead stems of *Bidens pilosa*. **Sexual morph**: Undetermined. **Asexual morph**: *Conidiomata* 100–150 \times 110–150 μ m (\bar{x} = 111 \times 130 μ m, n = 5), pycnidial, solitary, immersed to semi-immersed, uniloculate, globose, yellowish brown to brown, ostiolate. *Peridium* 13–20 μ m wide, comprising 1–2 layers of brown cells of *textura angularis*. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* 1–2 \times 3–4 μ m (\bar{x} = 1.4 \times 3.6 μ m, n = 5) phialidic, hyaline. *Conidia* 5–11 \times 4–10 μ m (\bar{x} = 12.8 \times 6.1 μ m, n = 20), ovoid to obpyriform, yellowish brown to brown, aseptate when immature, becoming brown and 1-septate at maturity, thick-walled, verruculose.

Table 3. Synopsis of recorded asexual morph of *Chromolaenicola* species.

Species	Conidiomata (μ m)	Peridium (μ m)	Conidiogenous cells (μ m)	Conidia (μ m)	Host/substrate	Ref.
<i>C. ananasi</i> (MFLU 23-0167)	–	–	–	7–8 \times 4–5	<i>Ananas comosus</i> (Bromeliaceae)	[30]
<i>C. chiangraiensis</i> (MFLUCC 17-1493)	–	–	3.5–6.5 \times 1–2	9–14 \times 6–9	<i>Chromolaena odorata</i> (Asteraceae)	[17]
<i>C. chaingraiensis</i> (MFLUCC 24-0058)	–	–	2–4 \times 1–2	9–13 \times 6–10	<i>Bidens pilosa</i> (Asteraceae)	This study
<i>C. clematidis</i> (MFLUCC 17-2075)	76–145 \times 107–128	5–10	2.6–4.5 \times 4–7	7–10 \times 4.5–7	<i>Clematis subumbellata</i> (Ranunculaceae)	[29]
<i>C. lampangensis</i> (MFLUCC 17-1462)	150–230 \times 170–270	10–20	–	12–15 \times 4–6.5	<i>Chromolaena odorata</i> (Asteraceae)	[17]
<i>C. siamensis</i> (= <i>Cylindroaseptospora siamensis</i> , MFLUCC 17-2527)	110–165 \times 140–190	15–38	6.5–7.4 \times 3.2–4.7	7.2–9.4 \times 5.4–6.5	<i>Leucaena</i> sp. (Fabaceae)	[28]
<i>C. siamensis</i> (MFLUCC 24-0057)	130–235 \times 170–230	15–20	1–2.5 \times 2–4	7–15 \times 5–10	<i>Bidens pilosa</i> (Asteraceae)	This study
<i>C. thailandensis</i> (MFLUCC 24-0056)	100–150 \times 110–150	13–20	1–2 \times 3–4	5–11 \times 4–10	<i>Bidens pilosa</i> (Asteraceae)	This study




Fig. 3 *Chromolaenicola siamensis* (MFLU 24-0029, new host record). (a), (b) Conidiomata on the substrate. (c) Section through conidiomata. (d) Peridium. (e), (f) Conidiogenous cells. (g)–(l) Conidia. (m) Germinating conidia. (n) Culture on MEA. Scale bars: (a), (b) = 500 μ m, (c) = 50 μ m, (d) = 20 μ m, (e)–(m) = 5 μ m, (n) = 10 mm.




Fig. 4 *Chromolaenicola thailandensis* (MFLU 24-0028, new host record, first report of asexual morph). (a), (b) Colonies on substrate. (c) Section through conidiomata. (d) Peridium. (e) Conidiogenous cells. (f)–(h) Conidia. (i) Germinating conidia. (j) Culture on MEA. Scale bars: (a), (b) = 500 μ m, (c) = 50 μ m, (d)–(i) = 10 μ m, (j) = 10 mm.

Culture characteristics: Conidia germinating on MEA within 24 h, 21 mm after 10 d at room temperature, irregular, entire margin, smooth, wrinkled, pale yellow on the surface, curled and brown in reverse (Fig. 5).

Material examined: Thailand, Chiang Rai Province, Doi Pui, on dead stems of *Bidens pilosa*, 14 March 2023, Zin Hnin Htet (BP-DP-2, MFLU 24-0028, **new asexual morph record**); living culture MFLUCC 24-0056.

Notes: *Chromolaenicola thailandensis* (MFLUCC 17-1510, MFLUCC 17-1475) was found in its sexual morph in nature (Table 4)^[17]. In the current study, we collected an asexual morph of *C. thailandensis* (MFLUCC 24-0056) on the dead stems of *Bidens pilosa*. However, we could not obtain its sexual morph in culture; hence failed to compare its morphology with *C. thailandensis* (MFLUCC 17-1510, MFLUCC 17-1475). Furthermore, there are no significant base pair differences in all five gene regions between our strain (MFLUCC 24-0056) and *C. thailandensis* (MFLUCC 17-1510). Therefore, we reported our strains as the new asexual morph of *C. thailandensis* and also the new host record from *Bidens pilosa* (Asteraceae) while previous strains of *C. thailandensis* was recorded on *Chromolaena odorata* (Asteraceae) in Chiang Rai Province, Thailand^[17].

Preliminary screening for antibacterial activity

In this study, we observed the antibacterial activity of our newly isolated *Chromolaenicola* species against *Bacillus subtilis*, *Escherichia coli*, and *Staphylococcus aureus*. Among the three isolates, MFLUCC 24-0056, MFLUCC 24-0057, and MFLUCC 24-0058 showed antibacterial activity against *Bacillus subtilis* (17, 18, and 10 mm, respectively) and considered partial inhibition compared to the positive control. However, they did not inhibit the growth of *E. coli* and *Staphylococcus aureus*. Moreover, we also re-confirmed the preliminary antimicrobial activity result of *Chromolaenicola* species previously studied (Table 5).

Discussion

Chromolaenicola species are found as saprobes in terrestrial on different plant hosts^[17,27–29]. In previous studies, seven *Chromolaenicola* species were reported from Thailand, and two from China^[17,27–30]. We also provided detailed information on all *Chromolaenicola* species (Tables 3 & 4). Based on the morphological and phylogenetic analyses, three new host records and one new asexual morph report is described in this paper. Although the conidial color and conidiogenous cell sizes of the *Chromolaenicola chiangraiensis* strains MFLUCC 24-0058 and MFLUCC 17-1493 are different, the base pair difference is not significant, and the latter was reported on *Chromolaena odorata* (Asteraceae). Therefore, we described our strain as a new host record on *Bidens pilosa* (Asteraceae). Furthermore, *Chromolaenicola siamensis* (MFLUCC 17-1527) was recorded on *Leucaena* sp. (Fabaceae) and herein we reported our strain (MFLUCC 24-0057) as a new host record on *Bidens pilosa* (Asteraceae). This record also marks the first occurrence of *C. siamensis* on Asteraceae species. Moreover, we found the asexual morph of *Chromolaenicola thailandensis* (MFLUCC 24-0056) for the first time in our study, and this is also the new host record on *Bidens pilosa* (Asteraceae).

In this study, we provided the preliminary screening results of all *Chromolaenicola* species, including known previous studies. Three isolates of *Chromolaenicola* species from northern Thailand underwent preliminary screening for antibacterial activity, and all three isolates showed partial inhibition of the growth of *Bacillus subtilis*. (Table 5). Previously, four *Chromolaenicola* species, *C. chaingraiensis* (MFLUCC 17-1493), *C. lampangensis* (MFLUCC 17-1462), *C. nanensis* (MFLUCC 17-1473), and *C. thailandensis* (MFLUCC 17-1510) have been studied for their potential antimicrobial properties^[17]. Among them, *C. lampangensis* (MFLUCC 17-1462), *C. nanensis* (MFLUCC




Fig. 5 Culture characteristics on MEA. (a) *Chromolaenicola thailandensis* (MFLUCC 24-0056), (b) *Chromolaenicola siamensis* (MFLUCC 24-0057), (c) *Chromolaenicola chiangraiensis* (MFLUCC 24-0058).

Table 4. Synopsis of recorded sexual morph of *Chromolaenicola* species.

Species	Ascomata (μm)	Peridium (μm)	Asci (μm)	Ascospores (μm)	Host/substrate	Ref.
<i>C. nanensis</i> (MFLUCC 17-1473)	210–230 × 200–220	15–20	110–145 × 10–12.5	16–20 × 7.5–9	<i>Chromolaena odorata</i> (Asteraceae)	[17]
<i>C. spindi</i> (KUMCC 21-0564)	420–530 × 270–350	15–25	125–155 × 12–16	16–23 × 6.5–9.5	<i>Sapindus rarak</i> (Spindaceae)	[27]
<i>C. thailandensis</i> (MFLUCC 17-1510)	145–225 × 175–240	10–20	90–160 × 10–14	16–24 × 9–11	<i>Chromolaena odorata</i> (Asteraceae)	[17]

Table 5. Preliminary antimicrobial activity result of *Chromolaenicola* species.

Species	Zone of inhibition (mm); Ampicillin (+)				Ref.
	<i>Bacillus subtilis</i>	<i>Escherichia coli</i>	<i>Staphylococcus aureus</i>	<i>Mucor plumbeus</i>	
<i>Chromolaenicola ananasi</i> (MFLU 23-0167)	N/A	N/A	N/A	N/A	[30]
<i>C. chaingraiensis</i> (MFLUCC 24-0058)	10 mm inhibition	no inhibition	No inhibition	N/A	This study
<i>C. chiangraiensis</i> (MFLUCC 17-1493)	No inhibition	No inhibition	N/A	no inhibition	[17]
<i>C. clematidis</i> (MFLUCC 17-2075)	N/A	N/A	N/A	N/A	[29]
<i>C. lampangensis</i> (MFLUCC 17-1462)	No inhibition	no inhibition	N/A	14 mm inhibition	[17]
<i>C. nanensis</i> (MFLUCC 17-1473)	No inhibition	No inhibition	N/A	12 mm inhibition	[17]
<i>C. siamensis</i> (MFLUCC 17-2527)	N/A	N/A	N/A	N/A	[28]
<i>C. siamensis</i> (MFLUCC 24-0057)	18 mm inhibition	no inhibition	no inhibition	N/A	This study
<i>C. spindii</i> (KUMCC 21-0564)	N/A	N/A	N/A	N/A	[27]
<i>C. thailandensis</i> (MFLUCC 17-1510)	No inhibition	No inhibition	N/A	No inhibition	[17]
<i>C. thailandensis</i> (MFLUCC 24-0056)	17 mm inhibition	No inhibition	No inhibition	N/A	This study

N/A: Not available; Positive control (+): Ampicillin.

17-1473) can inhibit the growth of *Mucor plumbeus*. The rest of the species of this genus have not yet been explored for their potential biological properties^[27–30]. In our study, the agar plug diffusion method was used to diffuse an antimicrobial drug into the agar, inhibiting bacterial growth, and then assessing the inhibition zone^[17,57]. Several benefits of the agar plug diffusion method include its inexpensive cost and ability to examine a large number of fungal cultures quickly^[57]. However, because it is impossible to determine the amount of antimicrobial agent diffused into the agar medium, this method is not appropriate for determining the minimum inhibitory concentration (MIC)^[57]. Our study will contribute to the knowledge of the species diversity in *Chromolaenicola* and insight into their potential biological properties, which will be useful for further research.

Author contributions

The authors confirm contributions to the paper as follows: Fungal specimen collection and isolation, and manuscript writing: Htet ZH; fungal identification and contributed in the revision of the manuscript: Mapook A, Chethana KWT. All authors have read and agreed to the published version of the manuscript. All authors reviewed the results and approved the final version of the manuscript.

Data availability

The data generated and analyzed during this study are available in the article. DNA sequence data are available in the GenBank database, and the accession numbers are provided in Table 1.

Acknowledgments

We would like to express our gratitude to the Center of Excellence in Fungal Research, as well as the Basic Research Fund support from the National Science, Research and Innovation Fund (Grant No. 652A01001), entitled 'Studies of fungi associated with Asteraceae and the discovery of biological properties'. Zin Hnin Htet would like to thank Mae Fah Luang University for giving her a scholarship to pursue a PhD and the re-inventing grant provided by the university for the molecular work 'Mae Fah Luang University Grant - Reinventing University System' (Grant No. 652A16049). Ausana Mapook thanks Mae

Fah Luang University for funding 'Taxonomy, Phylogeny, Risk Assessment, and Potential Impact of Fungi on Siam Weed in Northern Thailand' (Grant No. 651A16029).

Conflict of interest

The authors declare that they have no conflict of interest.

Dates

Received 18 March 2024; Revised 22 April 2024; Accepted 2 May 2024; Published online 22 May 2024

References

- Hooper DU, Chapin FS III, Ewel JJ, Hector A, Inchausti P, et al. 2005. Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecological Monographs* 75(1):3–35
- Hejda M, Pyšek P, Jarošík V. 2009. Impact of invasive plants on the species richness, diversity and composition of invaded communities. *Journal of Ecology* 97(3):393–403
- Pyšek P, Jarošík V, Hulme PE, Pergl J, Hejda M, et al. 2012. A global assessment of invasive plant impacts on resident species, communities and ecosystems: the interaction of impact measures, invading species' traits and environment. *Global Change Biology* 18(5):1725–37
- Blackburn TM, Essl F, Evans T, Hulme PE, Jeschke JM, et al. 2014. A unified classification of alien species based on the magnitude of their environmental impacts. *PLoS Biology* 12(5):e1001850
- Jeschke JM, Bacher S, Blackburn TM, Dick JTA, Essl F, et al. 2014. Defining the impact of non-native species. *Conservation Biology* 28(5):1188–94
- Kumschick S, Bacher S, Evans T, Marková Z, Pergl J, et al. 2015. Comparing impacts of alien plants and animals in Europe using a standard scoring system. *Journal of Applied Ecology* 52(3):552–61
- Weidlich EWA, Flórido FG, Sorriani TB, Brancalion PH. 2020. Controlling invasive plant species in ecological restoration: A global review. *Journal of Applied Ecology* 57(9):1806–17
- D'Antonio CM, August-Schmidt E, Fernandez-Goig B. 2016. Invasive species and restoration challenges. In *Foundations of restoration ecology*, eds. Palmer MA, Zedler JB, Falk DA. Washington, D.C.: Island Press. pp. 216–44. https://doi.org/10.5822/978-1-61091-698-1_8
- Arthur GD, Naidoo KK, Coopoomsamy RM. 2012. *Bidens pilosa* L. : Agricultural and pharmaceutical importance. *Journal of Medicinal Plants Research* 6(17):3282–81
- Zungontiporn S. 2007. Some characteristics of *Bidens pilosa* L. var. *radiata* Scheff., a new invasive species in Thailand. *Proceeding of the 21st Asian Pacific Weed Science Society (APWSS) Conference*, 2–6

- October 2007, Peradeniya, Sri Lanka. Sri Lanka: Asian Pacific Weed Science Society.
11. Guatimosim E, Pinto HJ, Pereira OL, Fuga CAG, Vieira BS, et al. 2015. Pathogenic mycobiota of the weeds *Bidens pilosa* and *Bidens subalternans*. *Tropical Plant Pathology* 40:298–317
 12. Zhang F, Li Q, Yeger EH, Chen X, Shi Q, et al. 2018. AM fungi facilitate the competitive growth of two invasive plant species, *Ambrosia artemisiifolia* and *Bidens pilosa*. *Mycorrhiza* 28:703–15
 13. Li J, Jeewon R, Mortimer PE, Doilom M, Phookamsak R, et al. 2020. Multigene phylogeny and taxonomy of *Dendryphion hydei* and *Torula hydei* spp. nov. from herbaceous litter in northern Thailand. *PLoS One* 15(2):e0228067
 14. Abdou R, Scherlach K, Dahse HM, Sattler I, Hertweck C. 2010. Botryorhizines A–D, antifungal and cytotoxic depsidones from *Botryosphaeria rhodina*, an endophyte of the medicinal plant *Bidens pilosa*. *Phytochemistry* 71:110–16
 15. Munk A. 1953. The system of the Pyrenomycetes: A contribution to a natural classification of the group Sphaeriales sensu Lindau. *Dansk Botanisk Arkiv* 15:1–163
 16. Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, et al. 2022. Outline of Fungi and fungus-like taxa–2021. *Mycosphere* 13(1):53–453
 17. Mapook A, Hyde KD, McKenzie EHC, Jones EBG, Bhat DJ, et al. 2020. Taxonomic and phylogenetic contributions to fungi associated with the invasive weed *Chromolaena odorata* (Siam weed). *Fungal Diversity* 101:1–175
 18. Samarakoon BC, Wanasinghe DN, Samarakoon MC, Phookamsak R, McKenzie EHC, et al. 2020a. Multi-gene phylogenetic evidence suggests *Dictyoarthrinium* belongs in Didymosphaeriaceae (Pleosporales, Dothideomycetes) and *Dictyoarthrinium musae* sp. nov. on *Musa* from Thailand. *Mycosphaerella* 71:101–18
 19. Samarakoon BC, Phookamsak R, Wanasinghe DN, Chomnunti P, Hyde KD, et al. 2020b. Taxonomy and phylogenetic appraisal of *Spegazzinia musae* sp. nov. and *S. deightonii* (Didymosphaeriaceae, Pleosporales) on Musaceae from Thailand. *Mycosphaerella* 70:19–37
 20. Yuan Z, Druzhinina IS, Wang X, Zhang X, Peng L, et al. 2020. Insight into a highly polymorphic endophyte isolated from the roots of the halophytic seepweed *Suaeda salsa*: *Laburnicola rhizohalophila* sp. nov. (Didymosphaeriaceae, Pleosporales). *Fungal Biology* 124:327–37
 21. Dissanayake LS, Wijayawardene NN, Samarakoon MC, Hyde KD, Kang JC. 2021. The taxonomy and phylogeny of *Austropleospora ochracea* sp. nov. (Didymosphaeriaceae) from Guizhou, China. *Phytotaxa* 491:217–29
 22. Wong Chin JM, Puchooa D, Bahorun T, Jeewon R. 2021. Molecular characterization of marine fungi associated with *Haliclona* sp. (sponge) and *Turbinaria conoides* and *Sargassum portierianum* (brown algae). *Proc. of the National Academy of Sciences, India Section B: Biological Sciences* 91:643–56
 23. Suwannarach N, Kumla J, Lumyong S. 2021. *Spegazzinia camelliae* sp. nov. (Didymosphaeriaceae, Pleosporales), a new endophytic fungus from northern Thailand. *Phytotaxa* 483:117–28
 24. Sun YR, Zhang JY, Hyde KD, Wang Y, Jayawardena RS. 2023. Morphology and phylogeny reveal three *Montagnula* species from China and Thailand. *Plants* 12(4):738
 25. Index Fungorum. www.indexfungorum.org (Accessed on December 2023).
 26. Species Fungorum 2023. www.indexfungorum.org/names/Names.asp (Accessed on December 2023).
 27. Ren G, Wanasinghe DN, de Farias ARG, Hyde KD, Yasanthika E, et al. 2022. Taxonomic novelties of woody litter fungi (Didymosphaeriaceae, Pleosporales) from the Greater Mekong Subregion. *Biology* 11(11):1660
 28. Jayasiri SC, Hyde KD, Jones EBG, McKenzie EHC, Jeewon R, et al. 2019. Diversity, morphology and molecular phylogeny of Dothideomycetes on decaying wild seed pods and fruits. *Mycosphere* 10:1–186
 29. Phukhamsakda C, McKenzie EHC, Phillips AJL, Gareth Jones EB, Jayarama Bhat D, et al. 2020. Microfungi associated with *Clematis* (Ranunculaceae) with an integrated approach to delimiting species boundaries. *Fungal Diversity* 102:1–203
 30. Tian XG, Bao DF, Karunarathna SC, Jayawardena RS, Hyde KD, Bhat DJ, et al. 2024. Taxonomy and phylogeny of ascomycetes associated with selected economically important monocotyledons in China and Thailand. *Mycosphere* 15:1–274
 31. Senanayake IC, Rathnayaka AR, Marasinghe DS, Calabon MS, Gentekaki E, et al. 2020. Morphological approaches in studying fungi: Collection, examination, isolation, sporulation and preservation. *Mycosphere* 11:2678–754
 32. Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat J, Buyck B, et al. 2015. The Faces of Fungi database: fungal names linked with morphology, phylogeny and human impacts. *Fungal Diversity* 74:3–18
 33. Chaiwan N, Gomdola D, Wang S, Monkai J, Tibpromma S, et al. 2021. <https://gmsmicrofungi.org>: An online database providing updated information of microfungi in the Greater Mekong Subregion. *Mycosphere* 12:1513–26
 34. Mapook A, Boonmee S, Ariyawansa HA, Tibpromma S, Campesori E, et al. 2016. Taxonomic and phylogenetic placement of *Nodulosphaeria*. *Mycological Progress* 15:34
 35. Vilgaly R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172:4238–46
 36. White TJ, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In *PCR protocols: a guide to methods and applications*, eds. Innis MA, Gelfand DH, Sninsky JJ, White TJ. London, UK: Academic Press. pp. 315–22. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
 37. Rehner S. 2001. *Primers for Elongation Factor 1- α (EF1- α)*. Insect Biocontrol Laboratory: USDA, ARS, PSI.
 38. Liu YJ, Whelen S, Hall BD. 1999. Phylogenetic relationships among ascomycetes: evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16:1799–808
 39. Wijesinghe SN, Wanasinghe DN, Maharachchikumbura SSN, Wang Y, Al-Sadi AM, et al. 2020. *Bimuria omanensis* sp. nov. (Didymosphaeriaceae, Pleosporales) from Oman. *Phytotaxa* 449(2):97–108
 40. Vu D, Groenewald M, De Vries M, Gehrman T, Stielow B, et al. 2019. Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Studies in mycology* 92(1):135–54
 41. Ariyawansa HA, Tanaka K, Thambugala KM, Phookamsak R, Tian Q, et al. 2014. A molecular phylogenetic reappraisal of the *Didymosphaeriaceae* (= *Montagnulaceae*). *Fungal Diversity* 68:69–104
 42. Alidadi A, Javan-Nikkhah M, Kowsari M, Karami S, Rastaghi ME. 2018. Some species of fungi associated with declined Persian oak trees in Ilam province with emphasis on new records to mycobiota of Iran. *Rostaniha* 19:75–91
 43. Lumbsch HT, Lindemuth R. 2001. Major lineages of *Dothideomycetes* (Ascomycota) inferred from SSU and LSU rDNA sequences. *Mycological Research* 105(8):901–8
 44. Tennakoon DS, Thambugala KM, de Silva NI, Suwannarach N, Lumyong S. 2022. A taxonomic assessment of novel and remarkable fungal species in *Didymosphaeriaceae* (Pleosporales, Dothideomycetes) from plant litter. *Frontiers in Microbiology* 13:1016285
 45. Crous PW, Summerell BA, Shivas RG, Burgess TI, Decock CA, et al. 2012. Fungal Planet description sheets: 107–127. *Persoonia - Molecular Phylogeny and Evolution of Fungi* 28(1):138–82
 46. Tanaka K, Hirayama K, Yonezawa H, Sato G, Toriyabe A, et al. 2015. Revision of the *Massariaceae* (Pleosporales, Dothideomycetes). *Studies in Mycology* 82:75–136
 47. Swindell SR, Plasterer TN. 1997. SEQMAN. In *Sequence data analysis guidebook*, ed. Swindell SR. Totowa, NJ: Springer. pp. 75–89. <https://doi.org/10.1385/0-89603-358-9.75>

48. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30(12):2725–29
49. Rannala B, Yang Z. 1996. Probability distribution of molecular evolutionary trees: a new method of phylogenetic inference. *Journal of Molecular Evolution* 43:304–11
50. Lemmon AR, Brown JM, Stanger-Hall K, Lemmon EM. 2009. The effect of ambiguous data on phylogenetic estimates obtained by maximum likelihood and Bayesian inference. *Systematic Biology* 58:130–45
51. Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. 2010 Gateway Computing Environments Workshop (GCE), New Orleans, LA, USA, 14 November 2010. USA: IEEE. pp. 1–8. <https://doi.org/10.1109/GCE.2010.5676129>.
52. Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30(9):1312–13
53. Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, et al. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61:539–42
54. Zhaxybayeva O, Gogarten JP. 2002. Bootstrap, Bayesian probability and maximum likelihood mapping: exploring new tools for comparative genome analyses. *BMC Genomics* 3:1–15
55. Rambaut A. 2012. FigTree v. 1.40. University of Oxford.
56. Alam ST, Le TAN, Park JS, Kwon HC, Kang K. 2019. Antimicrobial biophotonic treatment of ampicillin-resistant *Pseudomonas aeruginosa* with hypericin and ampicillin cotreatment followed by orange light. *Pharmaceutics* 11:641
57. Balouiri M, Sadiki M, Ibnsouda SK. 2016. Methods for in vitro evaluating antimicrobial activity: A review. *Journal of Pharmaceutical Analysis* 6:71–79



Copyright: © 2024 by the author(s). Published by Maximum Academic Press, Fayetteville, GA. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.