

Comparative phosphoproteome analyses provide new insight into ethephon stimulated latex yield in rubber tree

Authors

Shuguang Yang[#], Xiaoyu Du[#],
Xiaomin Deng, Weimin Tian,
Jinquan Chao^{*}


Correspondence

tianwang208@163.com

In Brief

Ethephon stimulates latex production through the phosphorylation glycolysis and dephosphorylated cell wall metabolism.

Graphical abstract



Highlights

- 2,380 phosphoproteins are identified in latex by using Tandem Mass Tag (TMT) approach.
- 389 phosphoproteins show differential enrichment upon ethephon treatment.
- The phosphorylation of glycolysis and the dephosphorylation of cell wall metabolism contribute to ethephon stimulated latex production in rubber tree.

Citation: Yang S, Du X, Deng X, Tian W, Chao J. 2024. Comparative phosphoproteome analyses provide new insight into ethephon stimulated latex yield in rubber tree. *Tropical Plants* 3: e001 <https://doi.org/10.48130/tp-0024-0002>

Comparative phosphoproteome analyses provide new insight into ethephon stimulated latex yield in rubber tree

Shuguang Yang^{1#}, Xiaoyu Du^{1,2#}, Xiaomin Deng¹, Weimin Tian¹ and Jinquan Chao^{1,3*}

¹ Ministry of Agriculture and Rural Affairs Key Laboratory of Biology and Genetic Resources of Rubber Tree, State Key Laboratory Breeding Base of Cultivation and Physiology for Tropical Crops, Rubber Research Institute, Chinese Academy of Tropical Agricultural Sciences, Haikou 571101, China

² College of Advance Agricultural Sciences, Zhejiang A&F University, Lin'an 311300, China

³ National Key Laboratory for Tropical Crop Breeding, Sanya Research Institute, Chinese Academy of Tropical Agricultural Sciences, Sanya 572024, China

Authors contributed equally: Shuguang Yang, Xiaoyu Du

* Corresponding author, E-mail: tianwang208@163.com

Abstract

Phosphorylation is one of the important post translational modifications that influences protein activity, expression, and interactions. Natural rubber (NR) is an important industrial raw material, mainly obtained from rubber tree. In production, the application of ethephon can effectively increase NR yield. In the present study, we investigated the phosphoproteomic profiling of latex upon ethephon treatment, by using the Tandem Mass Tag (TMT) approach. A total of 5,548 phosphorylation sites on 2,380 proteins were identified. Of which, 389 phosphoproteins showed differential enrichment under the threshold of fold change (FC) ≥ 1.3 (upregulated) or ≤ 0.769 (downregulated) and p -value ≤ 0.05 . Gene Ontology (GO) enrichment analysis showed pathways such as glycolysis and reactive oxygen species (ROS) scavenging were activated, while processes such as tricarboxylic acid (TCA) cycle, cell wall biosynthesis, and cytoskeleton formation were inhibited upon ethephon treatment. The results may provide novel insights into the mechanism underlying ethephon-induced rubber latex production in rubber tree.

Citation: Yang S, Du X, Deng X, Tian W, Chao J. 2024. Comparative phosphoproteome analyses provide new insight into ethephon stimulated latex yield in rubber tree. *Tropical Plants* 3: e001 <https://doi.org/10.48130/tp-0024-0002>

Introduction

Post translational modification (PTM) is one of the important ways to influence protein activity, expression, and interactions^[1, 2]. Over the past few decades, more than 400 types of PTMs have been discovered, including phosphorylation, methylation, acetylation, glycosylation, ubiquitination, etc.^[3]. Among them, phosphorylation is one of the most widely studied^[4, 5]. Phosphorylation is a reversible modification process. The phosphate group is added to the protein by a protein kinase and removed by a protein phosphatase^[6]. In eukaryotes, approximately one-third of proteins undergo phosphorylation modification^[7]. In the past, it was difficult to identify phosphorylated proteins. However, recently, large-scale protein phosphorylation analyses have been performed with the development of mass spectrometry technology. In wheat, Vu et al. identified 5,178 and 7,023 phosphorylated sites in leaf and spikelet, respectively^[8]. In *Lotus japonicus*, Zhang et al. identified 2,957 phosphopeptides during the various stages of nodule development^[9].

Natural rubber (NR) is vital to human society due to its excellent properties, including resilience, elasticity, abrasion, and impact resistance^[10]. More than 2,500 plants can produce NR, including *Eucommia ulmoides*, *Hevea brasiliensis*, *Taraxacum kok-saghyz*, *Parthenium argentatum*, etc^[11]. Among them, *Hevea brasiliensis* (termed as rubber tree) is the only commercialized species, providing more than 98% of the total NR production worldwide^[12]. The laticifers located in the inner bark of rubber tree are the sites for NR biosynthesis and storage^[13]. The latex is essentially the cytoplasm of laticifer cells, which contains 30%–40% NR. In production, latex is collected by regularly

tapping the rubber tree bark. As one of the vital phytohormones, ethylene plays an important role in regulating plant growth and development^[14,15]. In the rubber tree, application of ethephon (an ethylene releaser) can notably increase NR production per tapping by prolonging the duration of latex flow^[16]. The significant increase in rubber production has attracted the attention of many researchers, and a wealth of information have been obtained. At the transcriptional level, transcriptome analysis suggested that the genes related to glycolysis are up-regulated while the genes related to cell wall formation are down-regulated^[17]. At the protein level, proteins associated with sugar metabolism and rubber particle aggregation are influenced by ethephon treatment^[18,19]. In 2015, Wang et al. performed a proteomics analysis of ethylene-stimulated rubber latex and identified 59 differential phosphoprotein by using two-dimensional differential in-gel electrophoresis combined with Pro-Q Diamond dye^[20]. Here, we explore a high-resolution phosphoproteomic profiling of latex upon ethephon application by using the Tandem Mass Tag (TMT) approach, attempting to uncover the underlying mechanisms for the significant increase in latex production.

Materials and methods

Plant materials

Eighteen regularly tapped rubber trees of the *Hevea brasiliensis* clone CATAS7-33-97 were grown at the Experimental Farm of the Chinese Academy of Tropical Agricultural Sciences. To avoid the influence of endogenous hormones produced by tapping, all the selected trees were rested for 20 d. Of these,

Latex phosphoproteome upon ethephon treatment

nine rested trees were treated with 0.5% ethephon on the 1st, 3rd, and 5th days before tapping, and the remaining trees were used as control. Each treatment consisted of three individual trees as biological replicates. All the selected trees were tapped with a half spiral pattern, every 3 d (S/2, d/3). The latex samples were dropped into an equal volume of extracting buffer (0.4 M mannitol, 0.02 M Tris-HCl, 2 mM PMSF, 2 X phosphatase inhibitor cocktail), and subsequently stored at -80°C for protein extraction. The latex production was measured using a graduated cylinder after latex flow completely stopped, and the dry rubber content was determined according to the method published previously^[21].

Protein extraction and digestion

Latex proteins were extracted using a modified phenol/acetone extraction method^[22]. Briefly, the thawed latex samples were mixed with an equal volume of tris-saturated phenol. After shaking for 10 min, the samples were centrifuged at 18,000 rpm for 45 min at 4°C . The phenol phases were collected and then mixed in 4-fold volumes of ice-cold 10% (v/v) NH_4AC in acetone, and stored overnight at -20°C . The protein pellets were collected by centrifugation at 12,000 rpm for 20 min at 4°C and solubilized in 1 X PBS buffer with phosphatase inhibitor. The digestion was performed using trypsin with a ratio of 1:50 (w/w) at 37°C overnight. The peptide segments were desalinated with Strata X C18 and vacuum freeze-dried.

Tandem Mass Tag (TMT) labeling and phosphopeptide enrichment

The peptide segments were dissolved in 0.5 M TEAB and labeled according to the manufacturer of TMT Kit. The TiO_2 beads were added and slowly shaking for 2 h. The beads were collected by centrifugation at 5,000 rpm for 1 min and washed with washing buffer three times. Finally, the phosphopeptides were collected by eluted using elution buffer and vacuum freeze-dried.

Mass spectrometry analysis

The peptide segments were separated by an ultra-high performance liquid chromatography system and injected into an NSI ion source for ionization, followed by analysis by Q Active HF-X mass spectrometry. The ion source voltage was set to 2.0 kV, and the peptide parent ions and their secondary fragments were detected and analyzed using high-resolution Orbitrap. The scanning range of the primary mass spectrometry was set to 350–1,600 m/z, and the scanning resolution was set to 60,000. The scanning range of the secondary mass spectrometry was fixed at a starting point of 100 m/z, and the resolution of the secondary scanning was set to 30,000.

Phosphoproteins identification

The secondary mass spectrometry data were retrieved using Maxquant (v1.5.2.8) and searched against the rubber tree clone CATAS8-79 database^[23]. Trypsin was selected for the enzyme. The tolerance for mass error of the primary parent ion in First search and Main search was set to 20 ppm and 5 ppm, respectively. The tolerance for mass error of the secondary fragment ion was 0.02 Da. The false positive rate (FDR) for phosphoproteins identification was set to 1%. The differential enriched phosphoproteins (DEPs) were identified based on the threshold of fold change (FC) ≥ 1.3 (upregulated) or ≤ 0.769 (downregulated) and p -value ≤ 0.05 .

Bioinformatics analysis

For motif enrichment analysis, the sequence of 13 amino acids (including the modified sites and six upstream/downstream sites) were analyzed using MoMo software. For Gene Ontology (GO) analysis, the Blast2GO program was used to get GO annotation based on molecular function, biological process, and cellular component.

Statistical analysis

Statistical analysis was performed by GraphPad Prism (version 8.0) by analysis of variance (ANOVA) based on t-test. Standard error indicates three independent biological replicates. Asterisks indicate significant differences (* $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$).

Results

Physiological changes in latex production and dry rubber content upon ethephon treatment

We determined the latex production and dry rubber content of latex from three successive tappings treated with ethephon at different time points: 1 d, 3d and 5 d. Results showed that although there was no difference in rubber content of latex, there was a significant change in latex production (Fig. 1a, b). Compared to 1 d and 5 d, the latex production notably increased at 3 d after treatment, reached 4.12-fold, 6.51-fold, and 5.59-fold at three successive tappings. The latex of 3 d after treatment at tapping No.1 (as well as relative control) were selected for further phosphoproteomic analysis.

Overview of latex phosphoproteomic profiling

Using the TMT approach, a total of 5,548 phosphorylation sites originating from 2,380 proteins were identified (Fig. 2a). Among them, 87.53% (4,856/5,548) of the sites were from phosphoserine (pSer), 10.11% (561/5,548) were from phosphothreonine (pThr), 2.36% (131/5,548) were from phosphotyrosine (pTyr) (Fig. 2b). Among the 2,380 phosphoproteins, 1,210 (50.84%) contained only one phosphorylated site, while the remaining 1,170 (49.16%) contained more than two phosphorylated sites. An uncharacterized protein (encoded by *Hb07g045150.5*) had the most, with 33 phosphorylated sites.

Under the threshold of fold change (FC) ≥ 1.3 (upregulated) or ≤ 0.769 (downregulated) and p -value ≤ 0.05 , 389 differential enriched phosphoproteins (DEPs) were identified, including 246 up-regulation and 143 down-regulation respectively (Fig. 2c). Phosphorylation motifs enrichment analysis showed nine motifs enriched in up-regulation ([pSP], [pSPR], [SxxpS], [PxpSP], [pSPK], [LxKxpS], [RxxpS], [GpS], [KxxpS]), while three motifs enriched in up-regulation ([RxxpSP], [pSP], [SxxpS]) (Fig. 2d).

We then analyzed the enriched GO terms in both up-regulation and down-regulation (Fig. 3a, b). The numbers of terms for cell components, molecular functions and biological processes were 8, 8, and 14, respectively. For cell components, the terms enriched in up-regulation mainly related to DNA replication, such as 'nuclear chromosome', 'nuclear chromatin', 'chromatin', 'nuclear replication fork', 'replication fork protection complex', but it mainly related to cytoskeleton in down-regulation, such as 'actin cytoskeleton', 'cortical actin cytoskeleton', 'actin cortical patch', 'cortical cytoskeleton'. For molecular functions, the terms enriched in up-regulation mainly related to kinase and ethylene signal, such as 'nucleoside kinase activity', 'uridine




Fig. 1 (a) Latex production and (b) rubber content from ethephon-treated rubber trees. Significant difference was indicated by the asterisks above the bars (n.s. represents no significance, * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$). 1 d, 3 d, 5 d represent 0.5% ethephon treatment on the 1st, 3rd, and 5th days before tapping. Tapping No. represents tapping number.

kinase activity', 'kinase activity', 'protein serine/threonine kinase activity', 'ethylene binding', but it mainly related to cytoskeleton in down-regulation, such as 'actin-dependent ATPase activity', 'microfilament motor activity', 'actin binding', 'actin filament binding'. For biological processes, the terms enriched in up-regulation mainly related to secondary metabolism and carbohydrate metabolism, such as 'regulation of phenylpropanoid metabolic process', 'regulation of lignin biosynthetic process', 'glycosyl compound biosynthetic process', 'regulation of polysaccharide metabolic process', but it mainly related to development and cytoskeleton in down-regulation, such as 'regulation of root development', 'regulation of post-embryonic root development', 'regulation of post-embryonic development', 'regulation of stem cell division', 'regulation of seed development', 'actin cytoskeleton organization', 'cytoskeleton organization'.

DEPs related to carbohydrate metabolism and IPP pathway

In rubber tree, activated carbohydrate metabolism provides not only energy but also carbon skeletons for organic compound formation, including NR. Acetyl-CoA generated by glycolysis is the starting point of mevalonate (MVA) pathway, which produces isopentenyl pyrophosphate (IPP), the direct precursor of natural rubber. Here, we detected one sugar transporter protein (SWEET), two glycolysis related enzymes (PFK and F2KP), one MVA pathway rate-limiting enzyme (HMGR) that were up-regulated, and two tricarboxylic acid cycle related enzymes (CS and MDH) that were down-regulated (Fig. 4).

DEPs related to latex flow

Several factors, including turgor pressure caused by cell wall, agglutination formation, and the ROS scavenging system, can influence latex flow upon tapping. We detected three cell wall

related proteins (CESA1, CESA3, 4CL) and three agglutination formation related proteins (Myosin-15, Myosin-17, Gluc) that were down-regulated, while one ROS scavenging system member (SOD) was up-regulated (Fig. 5).

Protein kinases and protein phosphatases in DEPs

Kinase and phosphatase play important roles in dynamic phosphorylation. In this study, eight protein phosphatases (one PP2C, two tyrosine phosphatases, five Ser/Thr protein phosphatases) and 40 protein kinases (two CDPKs, two CIPKs, two MAPKs, one MAPKK, one MAPKKK, three CKs, two SnRKs, 27 Ser/Thr protein kinases) were detected in DEPs (Supplemental Table S1).

Discussion

High-resolution latex phosphoproteomic profiling upon ethephon treatment

Application of ethephon significantly increases latex production per tapping by prolonging the duration of latex flow in rubber tree. Research on the ethephon-enhanced latex production has mainly focused on transcriptomics or proteomics. Protein phosphorylation is one of the important PTM, playing a vital role in regulation of signal transduction, regulating cell proliferation, development, differentiation, and apoptosis. By using two-dimensional differential in-gel electrophoresis combined with Pro-Q Diamond dye, Wang et al. detect more than 200 protein spots and 59 phosphoproteins with differential enrichment upon ethylene treatment^[20]. In the present study, we identified 2,380 phosphoproteins by using the TMT approach. The number of phosphoproteins increased 10-folds compared to the previous study, suggesting the high throughput of TMT technology than that of 2-DE gel. Additionally, the

phosphorylated rubber elongation factor/small rubber particle proteins reported previously are not detected in this study. This may be attributed to the normal concentration (0.5%) ethephon used here, while the high concentration (3%) ethephon used by Wang et al.^[20].

Hundreds of DEPs upon ethephon treatment make it possible to globally analyze. Motif enrichment analysis of up-/down-regulation detected several conserved motifs. It is well-known that [pSP] is targeted by MAPKs and [RxxpS] is targeted by CDPK and SnRK^[24,25]. Here, we show [RxxpS] is one of the most enriched motifs in up-regulation, corresponding to the two CDPKs and two SnRKs detected in up-regulation. In addition, [pSP] is one of the most enriched motifs in both up-regulation and down-regulation, suggesting the complex MAPK signal network in latex of rubber tree.

Application of ethephon activates glycolysis and inhibits TCA cycle

Carbohydrate metabolism, such as sucrose metabolism and glycolysis, not only supplies energy but also generates a carbon skeleton for the formation of organic compounds including rubber. SWEET protein is a new class of sucrose transporters that function as bidirectional uniporters/facilitators and facilitate the diffusion of sucrose across cell membranes along a concentration gradient^[26]. In *Arabidopsis*, rapid phosphorylation of SWEET11/12 enhances the sucrose transport activity upon drought and abscisic acid treatments^[27]. Here, we notice that SWEET is phosphorylated upon ethephon treatment, speculating that it may be contributing to the sucrose transport into laticifer cell. Acetyl-CoA generated from glycolysis is an important carbon skeleton. It enters TCA cycle to produce energy or MVA pathway to produce IPP for rubber biosynthesis^[28]. PFK is one of the rate-limiting enzymes in glycolysis. Phosphorylation of PFK results in increased sensitivity to ATP inhibition and stronger binding of ATP to the inhibitory site of the enzyme^[29]. In the TCA cycle, oxaloacetate is the rate-limiting factor. It can produce citric acid by CS and be regenerated by MDH catalyzing malate^[30]. Here we show that PFK is up-regulated while both CS and MDH are down-regulated. HMGR is a key rate-limiting enzyme in the MVA pathway by controlling the carbon source entry^[31]. Here, we detect that it is up-regulated upon ethephon treatment. In summary, the activation of sucrose transport and glycolysis and the inhibition of TCA cycle ensure sufficient IPP for the MVA pathway.

Application of ethephon decreases the resistance of latex flow

The duration of latex flow is influenced by various factors, such as laticifer turgor pressure, plug formation at the end of severed laticifer, and latex redox^[32]. Turgor pressure is the initial driving force for latex flow after tapping. The properties of laticifer cell wall are closely related to turgor pressure. Lignin and cellulose are the two major components of cell wall. CESA is the key enzyme for cellulose biosynthesis. In *Arabidopsis*, CPK32 can interact with both CESA1 and CESA3, and specifically phosphorylates CESA3 on T672 residue *in vitro*. Overexpressing functionally defective CPK32 variant and phospho-dead mutation of CESA3 led to decreased motility of CSCs and reduced crystalline cellulose content in etiolated seedlings^[33]. Here we show CESA1 and CESA3, as well as 4-CL (a key enzyme

for lignin formation), reduce the level of phosphorylation, suggesting that ethephon treatment may increase turgor pressure by reducing lignin and cellulose content in laticifer. The formation and accumulation of protein-networks at the end of the laticifer's wounded site is the direct result of the cessation of latex flow after tapping^[34]. Cytoskeleton proteins, such as actin and myosin, play a key role in the formation of protein network. In *Physarum polycephalum*, phosphorylated myosin shows a significant increase in activity^[35]. Here we notice the reduction of phosphorylation level in both myosin-13 and myosin-15, hinting the formation of protein networks is affected upon ethephon treatment, which may contribute to the prolonging the duration of latex flow. The balance of ROS generation and scavenging in laticifer cells is an important factor for latex exploitation. High levels of ROS cause luteoids break, subsequently triggering the aggregation of latex^[36]. SOD is one of the important ROS scavenging enzymes. In 2001, Csar et al. reported that phosphorylation of Cu/Zn-SOD activates the G-CSF-mediated biological responses granulocyte^[37]. The increase in phosphorylation level of SOD upon ethephon treatment may reduce the level of ROS in latex, which benefits to latex exploitation after tapping. Finally, the application of ethephon decreases the resistance of latex flow caused by cell wall and protein network, resulting in the prolonging the duration of latex flow.

Conclusions

In this study, 2,380 phosphoproteins were identified in latex using TMT technology, and 389 phosphoproteins with differential enrichment were detected upon ethephon treatment. We speculate that the phosphorylation of key enzymes such as SWEET, F2KP, PFK, HMGR, SOD, while the dephosphorylation of key enzymes such as CS, MDH, 4CL, CESA, Myosin, Gluc, may play an important role in rubber yield increase upon ethephon treatment.

Author contributions

The authors confirm contribution to the paper as follows: study conception and design: Chao J, Tian W; ethephon treatment and latex collection: Yang S, Chao J; protein extraction: Chao J, Deng X; data analysis: Yang S, Du X; draft manuscript preparation: Chao J. All authors reviewed the results and approved the final version of the manuscript.

Data availability

The mass spectrometry phosphoproteomics data have been deposited to the Integrated Proteome Resources (iProX) with an accession: IPX0004047002. All data generated or analyzed during this study are included in this published article.

Acknowledgments

The research was supported by the National Natural Science Foundation of China (31700601), the Hainan Provincial Science and Technology Special Fund (ZDYF2022XDNY252), and the Earmarked Fund for China Agriculture Research System (CARS-




Fig. 3 Mass spectrometry analysis of phosphoproteins. (a) Statistical results of phosphoproteome identification. (b) Distribution of phosphorylated amino acids. (c) Differential enriched phosphoproteins identification. (d) Motif analysis of 246 upregulated phosphopeptides. (e) Motif analysis of 143 downregulated phosphopeptides.




Fig. 4 Protein phosphorylation related to carbohydrate metabolism and IPP pathway. The purple circles represent phosphorylation and the green circles represent dephosphorylation. PFK, phosphofructokinase; F2KP, fructose-2,6-bisphosphatase; CS, citrate synthase; MDH, malate dehydrogenase; HMGR, hydroxymethylglutaryl-CoA reductase.

33-YZ1).

Conflict of interest

The authors declare that they have no conflict of interest.

Jinquan Chao is the Editorial Board member of *Tropical Plants* who was blinded from reviewing or making decisions on the manuscript. The article was subject to the journal's standard procedures, with peer-review handled independently of this Editorial Board member and the research groups.




Fig. 5 Protein phosphorylation related to latex flow. The purple circle represents phosphorylation and the green circles represent dephosphorylation. CESA, Cellulose synthase, 4CL, 4-coumarate-CoA ligase; Gluc, glucan endo-1,3-beta-glucosidase; SOD, superoxide dismutase.

Supplementary information accompanies this paper at (<https://www.maxapress.com/article/doi/10.48130/tp-0024-0002>)

Dates

Received 16 November 2023; Revised 28 December 2023; Accepted 2 January 2024; Published online 15 January 2024

References

- Vu LD, Gevaert K, De Smet I. 2018. Protein language: Post-translational modifications talking to each other. *Trends In Plant Science* 23:1068–80
- Khan RA, Abbas N. 2023. Role of epigenetic and post-translational modifications in anthocyanin biosynthesis: A review. *Gene* 887:147694
- Han D, Yu Z, Lai J, Yang C. 2022. Post-translational modification: a strategic response to high temperature in plants. *abIOTECH* 15(3):49–64
- Ghelis T. 2011. Signal processing by protein tyrosine phosphorylation in plants. *Plant Signaling and Behavior* 6:942–51
- Silva-Sanchez C, Li H, Chen S. 2015. Recent advances and challenges in plant phosphoproteomics. *Proteomics* 15:1127–41
- Chao J, Huang Z, Yang S, Deng X, Tian W. 2020. Genome-wide identification and expression analysis of the phosphatase 2A family in rubber tree (*Hevea brasiliensis*). *PLoS One* 15:e0228219
- Dephoure N, Gould KL, Gygi SP, Kellogg DR. 2013. Mapping and analysis of phosphorylation sites: A quick guide for cell biologists. *Molecular Biology of The Cell* 24:535–42
- Vu LD, Stes E, Bel MV, Nelissen H, Maddelein D, et al. 2016. An up-to date workflow for plant (phospho) proteomics identifies differential drought-responsive, phosphorylation events in Maize leaves. *Journal of Proteome Research* 15:4304–17
- Zhang Z, Ke D, Hu M, Zhang C, Deng L, et al. 2019. Quantitative phosphoproteomic analyses provide evidence for extensive phosphorylation of regulatory proteins in the rhizobia-legume symbiosis. *Plant Molecular Biology* 100:265–83
- Hakimi NMF, Lee SH, Lum WC, Mohamad SF, Osman AI Edrus SS, et al. 2021. Surface Modified Nanocellulose and Its Reinforcement in Natural Rubber Matrix Nanocomposites: A Review. *Polymers* 13:3241
- Cherian S, Ryu SB, Cornish K. 2019. Natural rubber biosynthesis in plants, the rubber transferase complex, and metabolic engineering progress and prospects. *Plant Biotechnology Journal* 17:2041–61
- Priyadarshan PM, Goncalves PDS. 2003. *Hevea* gene pool for breeding. *Genetic Resources and Crop Evolution* 50:101–114
- Hao BZ, Wu JL. 2000. Laticifer differentiation in *Hevea brasiliensis*: induction by exogenous jasmonic acid and linolenic acid. *Annals of Botany* 85:37–43
- Wang KLC, Li H, Ecker JR. 2002. Ethylene biosynthesis and signaling networks. *The Plant Cell* 14:S131–S151
- Dubois M, Van den Broeck L, Inzé D. 2018. The pivotal role of ethylene in plant growth. *Trends In Plant Science* 23:311–23
- Nakano Y, Mitsuda N, Ide K, Mori T, Mira FR, et al. 2021. Transcriptome analysis of Pará rubber tree (*H. brasiliensis*) seedlings under ethylene stimulation. *BMC Plant Biology* 21:420

17. Liu JP, Zhuang YF, Guo XL, Li YJ. 2016. Molecular mechanism of ethylene stimulation of latex yield in rubber tree (*Hevea brasiliensis*) revealed by de novo sequencing and transcriptome analysis. *BMC Genomics* 17:257
18. Wang D, Xie Q, Sun Y, Tong Z, Chang L, et al. 2019. Proteomic Landscape Has Revealed Small Rubber Particles Are Crucial Rubber Biosynthetic Machines for Ethylene-Stimulation in Natural Rubber Production. *International Journal of Molecular Sciences* 20:5082
19. Dai L, Kang G, Nie Z, Li Y, Zeng R. 2016. Comparative proteomic analysis of latex from *Hevea brasiliensis* treated with Ethrel and methyl jasmonate using iTRAQ-coupled two-dimensional LC-MS/MS. *Journal of Proteomics* 132:167–75
20. Wang X, Wang D, Sun Y, Yang Q, Chang L, et al. 2015. Comprehensive Proteomics Analysis of Laticifer Latex Reveals New Insights into Ethylene Stimulation of Natural Rubber Production. *Scientific Reports* 5:13778
21. Chao J, Yang S, Chen Y, Tian WM. 2017. Transcript Profiling of *Hevea brasiliensis* during Latex Flow. *Frontiers in Plant Science* 8:1904
22. Wang X, Shi M, Lu X, Ma R, Wu C, et al. 2010. A method for protein extraction from different subcellular fractions of laticifer latex in *Hevea brasiliensis* compatible with 2-DE and MS. *Proteome Science* 8:35
23. Chao J, Wu S, Shi M, Xu X, Gao Q, et al. 2023. Genomic insight into domestication of rubber tree. *Nature Communications* 14:4651
24. Lee TY, Lin ZQ, Hsieh SJ, Bretaña NA, Lu CT. 2011. Exploiting maximal dependence decomposition to identify conserved motifs from a group of aligned signal sequences. *Bioinformatics* 27:1780–87
25. Amagai A, Honda Y, Ishikawa S, Hara Y, Kuwamura M, et al. 2018. Phosphoproteomic profiling reveals ABA-responsive phosphosignaling pathways in *Physcomitrella patens*. *The Plant Journal* 94:699–708
26. Ji J, Yang L, Fang Z, Zhang Y, Zhuang M, et al. 2022. Plant SWEET Family of Sugar Transporters: Structure, Evolution and Biological Functions. *Biomolecules* 12:205
27. Chen Q, Hu T, Li X, Song CP, Zhu JK, et al. 2022. Phosphorylation of SWEET sucrose transporters regulates plant root: shoot ratio under drought. *Nature Plants* 8:68–77
28. Tang C, Huang D, Yang J, Liu S, Sakr S, et al. 2010. The sucrose transporter *HsSUT3* plays an active role in sucrose loading to laticifer and rubber productivity in exploited trees of *Hevea brasiliensis* (para rubber tree). *Plant Cell and Environment* 33:1708–20
29. Kitajima S, Sakakibara R, Uyeda K. 1983. Significance of phosphorylation of phosphofructokinase. *Journal of Biological Chemistry* 258:13292–98
30. Ren M, Yang X, Bie J, Wang Z, Liu M, et al. 2020. Citrate synthase desuccinylation by SIRT5 promotes colon cancer cell proliferation and migration. *Biological Chemistry* 401:1031–1039
31. Wei H, Xu C, Movahedi A, Sun W, Li D, et al. 2019. Characterization and function of 3-hydroxy-3-methylglutaryl-CoA reductase in *Populus trichocarpa*: Overexpression of *PtHMGGR* enhances terpenoids in transgenic poplar. *Frontiers in Plant Science* 10:1476
32. Shi MJ, Cai FG, Tian WM. 2016. Ethrel-stimulated prolongation of latex flow in the rubber tree (*Hevea brasiliensis* Muell. Arg.): an Hev b 7-like protein acts as a universal antagonist of rubber particle aggregating factors from luteoids and C-serum. *Journal Of Biochemistry* 159:209–16
33. Xin X, Wei D, Lei L, Zheng H, Wallace IS, et al. 2023. CALCIUM-DEPENDENT PROTEIN KINASE32 regulates cellulose biosynthesis through post-translational modification of cellulose synthase. *New Phytologist* 239:2212–2224
34. Shi M, Li Y, Deng S, Wang D, Chen Y, et al. 2019. The formation and accumulation of protein-networks by physical interactions in the rapid occlusion of laticifer cells in rubber tree undergoing successive mechanical wounding. *BMC Plant Biology* 19:8
35. Kohama K, Kohno T, Okagaki T, Shimmen, T. 1991. Role of actin in the myosin-linked Ca^{2+} -regulation of ATP-dependent interaction between actin and myosin of a lower eukaryote, *Physarum polycephalum*. *Journal of Biochemistry* 110:508–513
36. Zhang Y, Leclercq J, Montoro P. 2017. Reactive oxygen species in *Hevea brasiliensis* latex and relevance to Tapping Panel Dryness. *Tree Physiology* 37:261–269
37. Csar XF, Wilson NJ, Strike P, Sparrow L, McMahon KA, et al. 2001. Copper/zinc superoxide dismutase is phosphorylated and modulated specifically by granulocyte-colony stimulating factor in myeloid cells. *Proteomics* 1:435–43



Copyright: © 2024 by the author(s). Published by Maximum Academic Press on behalf of Hainan University. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.